



POTSDAM-INSTITUT FÜR
KLIMAFOLGENFORSCHUNG

Originally published as:

Gleeson, T., Wang-Erlandsson, L., Porkka, M., Zipper, S. C., Jaramillo, F., [Gerten, D.](#), Fetzer, I., Cornell, S. E., Piemontese, L., Gordon, L. J., [Rockström, J.](#), Oki, T., Sivapalan, M., Wada, Y., Brauman, K. A., Flörke, M., Bierkens, M. F. P., Lehner, B., Keys, P., Kummu, M., Wagener, T., Dadson, S., Troy, T. J., Steffen, W., Falkenmark, M., Famiglietti, J. S. (2020): Illuminating water cycle modifications and Earth system resilience in the Anthropocene. - Water Resources Research, 56, 4, e2019WR024957.

DOI: <https://doi.org/10.1029/2019WR024957>

Water Resources Research

COMMISSIONED
MANUSCRIPT

10.1029/2019WR024957



Key Points:

- Earth system resilience depends on an improved understanding and management of water cycle modifications
- We identify four key functions of freshwater in the Earth system and evidence of regional to global regime shifts and disruptions
- The water planetary boundary is a compelling framework to improve our understanding and management of water cycle modifications in the Earth system

Correspondence to:

T. Gleeson,
tgleeson@uvic.ca

Citation:

Gleeson, T., Wang-Erlandsson, L., Porkka, M., Zipper, S. C., Jaramillo, F., Gerten, D., et al (2020). Illuminating water cycle modifications and Earth system resilience in the Anthropocene. *Water Resources Research*, 56, e2019WR024957. <https://doi.org/10.1029/2019WR024957>

Received 11 MAR 2019

Accepted 30 DEC 2019

Accepted article online 4 JAN 2020

Illuminating water cycle modifications and Earth system resilience in the Anthropocene

Tom Gleeson^{1,2}, Lan Wang-Erlandsson^{3,4,8}, Miina Porkka^{3,8}, Samuel C. Zipper^{1,2,5}, Fernando Jaramillo^{3,5}, Dieter Gerten^{6,7}, Ingo Fetzer^{3,8}, Sarah E. Cornell³, Luigi Piemontese³, Line J. Gordon³, Johan Rockström^{3,6}, Taikan Oki⁹, Murugesu Sivapalan^{10,11}, Yoshihide Wada^{12,15}, Kate A. Brauman¹³, Martina Flörke¹⁴, Marc F. P. Bierkens^{15,16}, Bernhard Lehner¹⁷, Patrick Keys¹⁸, Matti Kummu¹⁹, Thorsten Wagener^{20,21}, Simon Dadson²², Tara J. Troy¹, Will Steffen^{3,23}, Malin Falkenmark³, and James S. Famiglietti²⁴

¹Department of Civil Engineering, University of Victoria, Victoria, British Columbia, Canada, ²School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada, ³Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden, ⁴Research Institute for Humanity and Nature, Kyoto, Japan, ⁵Department of Physical Geography, Stockholm University, Stockholm, Sweden, ⁶Potsdam Institute for Climate Impact Research (PIK), Potsdam, Germany, ⁷Geography Department, Humboldt-Universität zu Berlin, Berlin, Germany, ⁸Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden, ⁹Integrated Research System for Sustainability Science, University of Tokyo, Tokyo, Japan, ¹⁰Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign, Urbana, IL, USA, ¹¹Department of Geography and Geographical Science, University of Illinois at Urbana-Champaign, Urbana, IL, USA, ¹²International Institute for Applied Systems Analysis, Laxenburg, Austria, ¹³Institute on the Environment, University of Minnesota, Saint Paul, MN, USA, ¹⁴Chair of Engineering Hydrology and Water Resources Management, Ruhr-University Bochum, Bochum, Germany, ¹⁵Physical Geography, Utrecht University, Utrecht, Netherlands, ¹⁶Deltares, Utrecht, Netherlands, ¹⁷Department of Geography, McGill University, Quebec, Canada, ¹⁸School of Global Environmental Sustainability, Colorado State University, Fort Collins, CO, USA, ¹⁹Water and Development Research Group, Aalto University, Espoo, Finland, ²⁰Department of Civil Engineering, University of Bristol, Bristol, UK, ²¹Cabot Institute, University of Bristol, Bristol, UK, ²²School of Geography and the Environment, University of Oxford, Oxford, UK, ²³Fenner School of Environment & Society, Australian National University, Canberra, Australian Capital Territory, Australia, ²⁴School of Environment and Sustainability and Global Institute for Water Security, University of Saskatchewan, Saskatoon, Saskatchewan, Canada, ²⁵Kansas Geological Survey, University of Kansas, Lawrence, KS, USA

Abstract Fresh water—the bloodstream of the biosphere—is at the center of the planetary drama of the Anthropocene. Water fluxes and stores regulate the Earth's climate and are essential for thriving aquatic and terrestrial ecosystems, as well as water, food, and energy security. But the water cycle is also being modified by humans at an unprecedented scale and rate. A holistic understanding of freshwater's role for Earth system resilience and the detection and monitoring of anthropogenic water cycle modifications across scales is urgent, yet existing methods and frameworks are not well suited for this. In this paper we highlight four core Earth system functions of water (hydroclimatic regulation, hydroecological regulation, storage, and transport) and key related processes. Building on systems and resilience theory, we review the evidence of regional-scale regime shifts and disruptions of the Earth system functions of water. We then propose a framework for detecting, monitoring, and establishing safe limits to water cycle modifications and identify four possible spatially explicit methods for their quantification. In sum, this paper presents an ambitious scientific and policy grand challenge that could substantially improve our understanding of the role of water in the Earth system and cross-scale management of water cycle modifications that would be a complementary approach to existing water management tools.

Plain language summary Freshwater is crucially important for all life on Earth. There is abundant research and evidence on how different processes within the water cycle regulate climate and support ecosystems, and by extension, human societies. Humans are also a major force disturbing those processes and modifying the water cycle. These modifications include, for instance, surface water withdrawals, groundwater pumping, deforestation and other land cover change, and ice melt due to warming climate. As most previous research on human–water interactions focuses on understanding systems at smaller scales, such as a watershed or a nation, comprehensive understanding of what human modifications of the water cycle mean for the stability of the planet is still lacking. In this paper we propose a new framework for analysing and establishing limits to a variety of human modifications of the water cycle,

to ensure that the stability of the Earth would not be compromised. We see this as an important and urgent scientific challenge that has the potential to substantially improve our understanding of the functioning of the Earth system and to inform local and global policy toward a more sustainable future.

1. Why is a Framework for Examining the Role of Water Cycle Modifications for Earth System Resilience Necessary?

Human pressure on fundamental planetary processes are pushing the Earth out of the Holocene geological epoch, the only period in Earth's history known to be capable of supporting sedentary, complex human civilisation. In this newly human shaped Anthropocene, the scale and magnitude of human drivers risk trigger critical transitions that jeopardize the habitability of Earth for human society (Barnosky et al., 2012; Steffen et al., 2018). Fresh water—the bloodstream of the biosphere—is at the center of this planetary drama: The water cycle is not only essential for myriad Earth system processes, interactions, and feedbacks, but it is also subject to anthropogenic manipulation at the global scale. Water, food, and energy security and sustainability depend on various water stores and fluxes that have been and are being modified by many and diverse processes, interactions, and feedbacks of global change in the Anthropocene.

Water flows regulate the Earth's climate system through mediation of the energy, carbon, and water balances and are a prerequisite for thriving aquatic and terrestrial ecosystems. Among others, soil moisture affects the Earth's albedo through supporting vegetation and contributing to cloud formation, carbon sequestration through regulation of biomass production, and moisture feedback on precipitation. Rivers transport half of the carbon sequestered by land to water bodies and respire half into the atmosphere (Biddanda, 2017). Critical ecosystem services require 90% of global evapotranspiration to function (Rockström et al., 1999), while streamflow sustains aquatic ecosystem functioning (Smakhtin 2004).

Holistically understanding, evaluating, and maintaining the water cycle's role for a resilient Earth system is extremely challenging and urgent in the Anthropocene, as the societal complexities interlock with the complex dynamics of the Earth system. Globally distributed and interconnected human activities have become the dominant force of modifications to the water cycle and already created local- to regional-scale water-mediated regime shifts. Research shows anthropogenic forcing to dominate changing river flows worldwide (Shiklomanov & Rodda, 2003), groundwater depletion (Aeschbach-Hertig & Gleeson, 2012; Bierkens & Wada, 2019), the partitioning of water on land (Jaramillo & Destouni, 2015a), as well as the spatial patterns and seasonal timing of evapotranspiration over continents (Gordon et al., 2005; Sterling et al., 2012). These water cycle modifications have had knock-on effects on critical Earth system functioning through modification of atmospheric moisture feedbacks (Wang-Erlandsson et al., 2018), tropical forest resilience (Hirota et al., 2011; Zemp et al., 2017; Staal et al., 2018), monsoon systems, and sea level (Wada et al., 2012). At the same time, the human impact on the water cycle is more internationally connected than ever, through flows of people, commodities, finance, technology, and information that enable emerging impacts on the water cycle through virtual water flows (Dalin et al., 2017; Hoekstra & Mekonnen, 2012; Oki et al., 2017; Porkka et al., 2012), land grabs (Rulli et al., 2013), and forest transition displacements (Meyfroidt et al., 2010).

The spatial heterogeneity and distribution of the water cycle, however, mean that water stores and fluxes are often analysed at local to regional scales of 10 to 1,000 s of km² (Archfield et al., 2015; Brown et al., 2015; Medema et al., 2008; Savenije et al., 2014). A textbook case of the need for a continental-to-global, rather than local-to-regional, perspective is the global interconnectedness between the Amazon Forest and the water cycle. Greenhouse gas emissions worldwide drive climate change that increases the frequency and severity of drought and fire in the Amazon (Aragão et al., 2018; Duffy et al., 2015), while global demand for agricultural products such as soybean and financial investments spur forest clearing (Barona et al., 2010; Galaz et al., 2018; Nepstad et al., 1994). Together, deforestation and climate change alter the regional rainfall patterns and shift the South American monsoon system and reverberate to impact on precipitation in the midlatitudes through teleconnections (D'Almeida et al., 2007; Lawrence and Vandecar, 2014; Nobre, 2014; Spera et al., 2016; Swann et al., 2015). Moreover, repeated severe droughts may ultimately undermine the Amazon forest's role as a global net carbon sink (Yang et al., 2018).

The scientific and ethical grand challenge of examining water cycle modifications for Earth system resilience is deeply connected to numerous research questions at the heart of current research frontiers and questions in hydrology: How do we understand and manage the interactions, dynamics, and connectivities in the global water system? (Alcamo et al., 2008; Bierkens et al., 2015; Fan et al., 2019; Vörösmarty et al., 2015; Wood et al., 2011). How do we consider humans as endogenous to the water system in the relatively newly emerging discipline of socio-hydrology? (Di Baldassarre et al., 2013.; Brown et al., 2015; Lund, 2015; Montanari et al., 2013; Sivapalan et al., 2012, 2014; Vogel et al., 2015; Wagener et al., 2010). How do we manage trade-offs between global development considered in the sustainable development goals (Bhaduri et al., 2016) and increasing pressure on the Earth system functioning of the global water cycle? How do existing water management and governance mechanisms and institutions respond to and influence global water cycling? How do interactions and feedbacks with the food and energy sectors impact the water cycle across scales? (Cai et al., 2018). Examining global water cycle modifications is a logical complement to these research challenges. To be able to navigate the planetary-scale dynamics of the water cycle and make connections to these other research questions, we first need to understand the following:

- What water-related changes may lead to supraregional or global tipping points—or more gradual yet equally detrimental transitions—related to water and Earth system functions?
- To what water-related changes, and in what regions, is the Earth system particularly vulnerable?
- How do local changes in stores and fluxes of water impact regional and global processes, and how do regional and global changes impact local processes?

In addition to these scientific questions, ideally, a framework should also be developed that recognizes all members of the global community as stakeholders of the global water cycle as a “global commons” and provides knowledge of globally distributed and aggregated limits to various water cycle modifications beyond which we may push the Earth system state into uncharted territory in terms of habitability for human civilisations.

Our objective is twofold: providing a planetary-scale overview of water’s role for maintaining Earth system functions and developing a framework for monitoring, detecting, and potentially acting on water cycle modifications. This work is based on multiple workshops, working groups, and intense collaboration and debate. Throughout, we focus on Earth system resilience using a series of concepts introduced in section 2 (dynamic systems, resilience theory, Earth system science, and the planetary boundary framework). Many manuscripts and textbooks have described the components and interactions of the stores and fluxes of the water cycle, but few (Falkenmark et al., 2019; Rockström et al., 2014) have highlighted the global water cycle as a dynamic, resilient system as we do in section 2. We then synthesize the functional role of water in the broader Earth system (section 3) and the evidence of regional regime shifts and disruptions of the Earth system functions of water (section 4). Finally, we propose a framework for defining global water cycle modifications using the planetary boundary framework (section 5) and conclude with a call to join us in this grand challenge (section 6). In the future, the methods for monitoring and detecting water cycle modifications could possibly be modified or adapted to examine other issues such as water scarcity, security, or virtual water. Herein “water” refers to terrestrial freshwater.

2. Earth System Resilience Theory

To motivate the following overview of resilience theory, we start with the potential role of human modifications of the water cycle in planetary-scale Earth system regime shift. Regime shifts are abrupt, persistent, and possibly irreversible changes in the system’s structure and function, which in general are well-documented across scales (Biggs et al., 2018; Rocha et al., 2015). Well-known local- to regional-scale regime shifts in hydrological systems include lake eutrophication even occurring in large lakes such as Lake Victoria (Hecky et al., 2010), lake depletion such as Aral Sea (Micklin, 2007; Micklin et al., 2016), and salinization such as in the Murray–Darling river basin (Overton et al., 2006). However, regime shifts occur also at the planetary scale. Known Earth system regime shifts include the last glacial–interglacial transition (Hoek, 2008), the “Big Five” mass extinctions (Barnosky et al., 2011) and the Cambrian explosion (Marshall, 2006). The evidence and understanding of past regime shifts in the Earth system is important, as current human pressure already exceeds the rate and magnitude of the forcings that precipitated the latest global-scale regime state shift, the last glacial–interglacial transition (Barnosky et al., 2012).

In general, regime shifts of a social–ecological system can occur through either a controlling parameter (such as rainfall) crossing a critical threshold leading to a “tipping” into a new state with internally stabilizing feedback processes or collapse through gradual change (Falkenmark et al., 2019). Regime shift through critical thresholds results in a state change that may be difficult or impossible to reverse due to the triggering of reinforcing positive feedbacks, whereas a regime shift through linear collapse of internal features or subsystems refers to a gradual system change that does not necessarily introduce significant differences in system feedbacks. Regime shift through threshold effects and reinforcing feedbacks can be illustrated through deforestation in the Amazon: Loss in moisture supply leads to rainfall reduction, which leads to forest loss below certain rainfall threshold, thus further rainfall reduction associated with increased risk of fire and further forest loss (Zemp et al., 2017). To restore the rainforest is more difficult because it involves hysteresis behavior enforced by internal feedbacks; in the Amazon, for example, counteracting self-amplifying feedbacks to restore a forested state requires higher rainfall levels than the threshold that originally induced forest loss. On the other hand, an example of linear collapse can be illustrated through river depletion, wherein principle, water levels can be restored by reversing the processes that led to their depletion (although related social–ecological systems such as fishing communities may exhibit hysteretic effects). To our best knowledge, there is no study comprehensively investigating whether human modifications of the water cycle have led, could be leading, or will lead to planetary-scale regime shifts in the Earth system.

The term resilience was originally introduced to refer to the capacity of a system to absorb or withstand disturbance that may precipitate a regime shift. This definition of resilience is sometimes referred to as persistence resilience; it draws originally from system dynamics understanding and is supported by a range of observation-based ecosystem studies (e.g., Holling, 1973; Scheffer et al., 2001; Walker et al., 2004). Persistence resilience can be used to describe all the examples of regime shifts described above and deals primarily with an agent-free environment (Donges & Barfuss, 2017) and therefore is also of primary relevance for understanding the biophysical aspects of Earth system resilience. Broader aspects of resilience of social–ecological systems (e.g., Folke et al., 2004; Holling, 1973; Holling & Gunderson, 2002) include the capacity of a system to adapt to changes and transform into a new desirable state. Complex interlinked social–ecological systems may undergo cycles of adaptation at different hierarchical levels, which are nested in the panarchy framework (Holling and Gunderson, 2002). At the global scale and highest level, social–ecological system dynamics have been addressed using world-system analysis (Chase-Dunn & Hall, 1997; Denmark et al., 2000; Gotts, 2007; Hall, 2000; Wallerstein, 1974) that to varying extent address socioeconomic feedbacks equally with biophysical feedbacks. The branch of Earth system science focusing on Earth system resilience to anthropogenic pressure that does not consider socioeconomic feedbacks as endogenous includes studies of tipping elements in the Earth system (Lenton et al., 2008; Steffen et al., 2018) and the planetary boundary framework (Rockström et al., 2009a, 2009b; Steffen, Richardson, et al., 2015).

Specifically, this paper focuses on the water cycle’s essential role for Earth system resilience, which is the Earth system’s ability to absorb or withstand perturbations and other stressors while essentially maintaining its structure and functions (see section 3). We recognize the three ways in which water interacts with resilience (Falkenmark et al., 2019; Rockström et al., 2014): (i) as a “source” of resilience, that is, through the generation of ecosystem services and functions in both terrestrial and aquatic systems; (ii) as a “victim” of change, for example, by being subject to land use change and pollution pressure; and (iii) as an “agent” of change, for example, by driving social shocks or vegetation change through modifications in the temporal or spatial distribution in the water cycle. By undermining resilience of local–regional systems, human modifications of the water cycle have already pushed many of them beyond collapse, such as through salinization, desertification, flow regulation, and eutrophication (see section 4 for evidence of water-related regime shifts).

In the Anthropocene, continued human pressure may now accumulate water cycle modifications to threats of planetary-scale tipping through four general pathways (Barnosky et al., 2012; Lenton et al., 2008; Rocha et al., 2018; Steffen et al., 2018): (1) extensive local-scale changes that trigger critical transitions over a large area; (2) global forcing that trigger local changes; (3) synergy, feedbacks, and cross-scale interactions through complex networks; and (4) tipping of major subsystems of the Earth system. The categorization of these pathways is idealized, and multiple mechanisms may concur in reality. The first pathway is supported by empirical patch-area and landscape-scale studies. For example, crossing certain thresholds of habitat loss or fragmentation may trigger abrupt, landscape-wide species extinction (Pardini et al., 2010). At

Earth system scale, it has been suggested that a 50% transformation of the Earth's terrestrial ecosystems may cause global-scale tipping to occur (Noss et al., 2012) and, for example, that 85% of tropical and boreal forests need to remain in order to safeguard the functioning of the Earth system (Rockström et al., 2009a, 2009b; Steffen, Richardson, et al., 2015). The second pathway can be illustrated by the role of Amazon as carbon sink, which if reverted to carbon source through drought-induced forest loss, may cause a global-scale forcing that triggers local-scale changes in the water cycle. The third pathway is supported by network analyses that show that regime shifts may interact through cascades and cross-scale interactions (e.g., Rocha et al., 2018). Dynamic and complex interactions and cascades have also raised the concern that processes that were previously thought to be confined to regional concern may in fact have planetary resilience implications (Rocha et al., 2018; Lenton et al., 2008, Steffen et al., 2018). Finally, the fourth pathway suggests that human pressure on tipping elements, which is the key subsystems of the Earth at subcontinental scale that can be switched into a different state following minor perturbations (Lenton et al., 2008), may have the ability to destabilize the current planetary state (Steffen et al., 2018).

We seek to understand the role of water cycle modifications for Earth system dynamics through the lens of the planetary boundary framework (section 5), as it is to date the most systematic effort for comprehensively addressing the question “What are the non-negotiable planetary preconditions that humanity needs to respect in order to avoid the risk of deleterious or even catastrophic environmental change at continental to global scales?” (Rockström et al., 2009b). Planetary boundaries are defined as biogeophysical boundaries for the processes and systems, which together regulate the state of the Earth system; the framework is not to be confused with the “planetary boundary layer” used in atmospheric science (Vilà-Guerau de Arellan et al., 2015). The planetary boundary framework is based on (i) identifying relevant biogeochemical processes that regulate the stability of the Earth system and (ii) determining the limit of human perturbation of these critical processes. Crossing any of the planetary boundaries could destabilize essential Earth system processes (Rockström et al., 2009a, 2009b; Steffen, Richardson, et al., 2015) and move the Earth away from Holocene conditions during which human societies developed and proliferated. Nine planetary boundary processes and systems have been identified (Rockström et al., 2009a, 2009b). For each boundary process/system, the planetary boundary should be defined based on the relationship between a control variable (a quantifiable, possibly spatially distributed biophysical indicator of main processes controlling the Earth system process in question, over which humans can exert some influence) and a response variable (an aspect of the Earth system that defines Earth's stable Holocene-like conditions and is affected by a change in the control variable), though several planetary boundaries do not yet have clearly defined control and/or response variables. It is important to note that the planetary boundary framework is based on biophysical resilience (the ability of a system to absorb or withstand disturbance and possibly ecosystems' ability to adapt to changing conditions) rather than the more socioecological resilience that includes societal adaptation, transformation, and panarchy.

The current planetary boundary for human freshwater use is based on a global sum of the average annual surface water flow in rivers relative to environmental flow requirements. Even if refined by complementary subglobal boundaries representing rivers' environmental flow requirements (Gerten et al., 2013; Steffen, Richardson, et al., 2015), this provisional planetary boundary and its current status have been critiqued (Heistermann, 2017; Jaramillo & Destouni, 2015b; Gleeson et al., 2019), as it does not reflect all types of human interference with the complex global water cycle and Earth system (Figure 1). Gleeson et al. (2019) suggest that a water planetary boundary would be more scientifically robust and more useful in decision-making frameworks if it was redesigned to consider more specifically how climate and living ecosystems respond to changes in the different forms of water on Earth: atmospheric water, soil moisture, groundwater and frozen water, as well as surface water. In section 5 we suggest how the planetary boundary framework can be used to better understand how water cycle modifications could potentially be impacting Earth system resilience, based on our following review.

3. The Earth System Functions of Water

The water cycle or hydrosphere is a complex system with different stores interacting with varying strengths and over a wide range of scales with other components of the Earth system such as atmosphere, biosphere, and lithosphere (Figure 1). Building on previous attempts in the systems and resilience literature




Figure 1. (a) The core functions of water in the Earth system. The five stores of the freshwater hydrosphere (colored circles in center), major components of the Earth system (outer ring), and detailed Earth system components underlying the different planetary boundaries (inner gray ring) are shown. The arrows denote processes linking the water stores and the Earth system components, color coded by water function (hydroclimate, hydroecology, storage, and transport). Since we focus on the near-surface hydrosphere, we consider land (part of the lithosphere) and ocean (part of the hydrosphere) as important related Earth system components. The interaction between stores are shown schematically but are not the focus since these are described in many hydrology textbooks. This diagram highlights the complex interactions between water stores and Earth system components more comprehensively than (b) common representations of the water cycle (modified from Oki & Kanae, 2006). Freshwater use is one of the current planetary boundaries yet affecting only a small component of (b) the hydrosphere and only representing a single function of water in the Earth system (see inset small circle and red text). Note that in figures, hydroclimatic and hydroecological regulation are shortened to hydroclimate and hydroecology. P, precipitation; ET, evapotranspiration.

(Rockström et al., 2014), seminal hydrology evaluations, reports, and textbooks (Dingman, 2002; National Research Council, 1991; Oki & Kanae, 2006; Tang & Oki, 2016; UNESCO, 1978) here we highlight four core Earth system functions of water: (1) hydroclimatic regulation, (2) hydroecological regulation, (3) storage, and (4) transport. These Earth system functions of water are different from watershed functions (Black, 1997; Wagener et al., 2007), which focus on hydrologic functions generally at smaller scales, not explicitly considering water in the broader Earth system. The Earth system functions of water are also different to water functions for social–ecological resilience in the Anthropocene (Falkenmark et al., 2019)

that distinguish green and blue water functions for social–ecological resilience, whereas we focus on the functions of water explicitly for Earth system stability, independent of green or blue origin. Inevitably, this description and related citations are nonexhaustive and serve primarily to outline a scientific foundation examining global water cycle modifications and for the water planetary boundary.

3.1. Hydroclimatic Regulation

Water exchange between the atmosphere, land surface, soil, ice and snow masses, oceans, and groundwater regulates the Earth's climate system through mediation of the energy, carbon, and water balance. Water vapor is the most powerful greenhouse gas due to its infrared absorption spectrum, heat storage capacity, and abundance in the atmosphere (Mitchell, 1989; Rodhe, 1990). Additionally, water vapor also forms clouds that reflect incoming solar radiation and absorb outgoing longwave radiation, with an overall effect on the Earth's energy balance that depends on cloud thickness, altitude, and constituent particles. Water vapor and the evaporation–condensation cycle is also the primary mechanism by which heat is redistributed from the equator to the poles (Henshaw et al., 2000). Soil moisture, surface water, and frozen water all directly or indirectly influence the albedo of the Earth's surface and thus the radiative balance. Soil moisture availability and surface water further affect carbon sinks and sources through mediating photosynthesis, oxygenation of soil, carbon transport, and carbon storage (Intergovernmental Panel on Climate Change, 2013). About half of the carbon sequestered by land is transported by rivers to water bodies, of which half is respired into the atmosphere (Biddanda, 2017). Finally, precipitation is influenced by evapotranspiration from land and soil moisture through boundary layer dynamics (Guillod et al., 2015), moisture recycling (van der Ent et al., 2010), and atmospheric circulation regulation (Tuinenburg, 2013). It should be noted that this core function has been undergoing major changes during the Anthropocene because of global climate change, which is often referred to as the “intensification” of the water cycle as reviewed by Huntington (2006), and a recent framework has been proposed for quantifying this intensification (Huntington et al., 2018).

3.2. Hydroecological Regulation

Water's hydroecological function enables and connects life on land and in aquatic ecosystems and creates and sustains the ecosystems that human societies and Earth system stability depend on (Gerten, 2013). This hydroecological function can be described by the quantity of water present at different times within the year relative to an ecosystem's water requirements. In aquatic ecosystems, this role of freshwater is often referred to as “environmental flows” (Acreman et al., 2014; Poff et al., 2009; Poff & Matthews, 2013). In terrestrial systems, the quantity and timing of available water relative to a species' physiological requirements is assigned as “hydrologic niche” and, along with other environmental constraints, drives species composition and ecosystem function (Booth & Loheide, 2012; Deane et al., 2017; Henszey et al., 2004). Changes to the quantity and timing of water availability can impact biosphere integrity and make ecosystems more vulnerable to drought or flooding and/or enable the invasion of non-native species (Catford et al., 2014; Pool et al., 2010; Zipper et al., 2017). Water's hydroecological functions are closely connected to its hydroclimatic and storage functions. Almost all water stored on land is of atmospheric origin, and surface waters (streams, lakes, and wetlands) harbor various types of aquatic ecosystems. Moreover, groundwater stores buffer aquatic ecosystems from the effects of short-term climatic variability. Hydroecological regulation is also closely tied to water's transport function as sediment and nutrient fluxes are critical determinants of aquatic habitat formation (Belmont & Foufoula-Georgiou, 2017; Motew et al., 2017; National Marine Fisheries Service, 2016).

3.3. Storage

Freshwater storage in groundwater, lakes, wetlands, reservoirs, and frozen water primarily interacts with the Earth system as a control over sea level. Globally, freshwater storage is dominated by frozen water in the polar ice sheets (Gleick, 2000). Mass loss due to ice melt is widespread and accelerating in both the Antarctic and Greenland ice sheets (Velicogna et al., 2014), and melt from the ice sheets increases the total volume of water in the oceans leading to sea level rise, exacerbated by thermal expansion of the oceans caused by global warming (Abraham et al., 2013). Groundwater is the second largest store of freshwater, and reductions in global groundwater storage due to groundwater pumping are a secondary contributor to global sea level (Wada et al., 2016), though the magnitude of this flux is dwarfed by the impacts of ice

melting (Reager et al., 2016). Increased storage in surface reservoirs plays only a minor role in total global water volume storage, yet it has the capacity to substantially distort river flow regimes, including the drying up of streams, and to reduce hydrologic connectivity (Grill et al., 2019). Loss of storage due to changes in lakes and wetlands, groundwater depletion, or reduced snowpack and/or mountain glaciers may also impact the Earth system via locally important alterations to the timing, magnitude, and temperature of streamflow (Dickerson-Lange & Mitchell, 2014; Gleeson & Richter, 2018; Immerzeel et al., 2010; Watson et al., 2014), which can have cascading effects on ecosystems and society (Xu et al., 2009).

3.4. Transport

The spatial and temporal dynamics of water are fundamental for moving, displacing, and diluting sediment and dissolved constituents including nutrients on the surface or within soils (Earle et al., 2015). Chemical weathering, mineral soil leaching, and transport of artificial fertilizers and chemicals into adjacent rivers, lakes and streams, and finally into the oceans (Earle et al., 2015; McGuire & McDonnell, 2006) impact water quality and biodiversity (Smith & Schindler, 2009). Water can either stabilize or destabilize landscapes (e.g., flooding) (Earle et al., 2015; Summerfield, 2005, 2014). Deposition of soil by water flux within and between these shape determine the function and geological shape of landscapes (Ellis et al., 2002; Wiens et al., 2005). Water and ice are responsible for a large amount of sediment transport on the surface of the Earth and are important in many geological processes such as rock and landform formation and erosion (Earle et al., 2015; Summerfield, 2005, 2014). Dilution of minerals and nutrients in soil additionally controls soil and above-ground biome characteristic (Ellis et al., 2002; Tölgyessy, 1993; Wiens et al., 2005).

4. Evidence of Regional Regime Shifts and Disruptions of the Earth System Functions of Water

Evidence of global water cycle modifications on Earth system functions is scattered, so here we highlight the existence and also the limitations of current knowledge of regional regime shifts and possible tipping elements for each water store. We point out key knowledge gaps that are essential to examine in the process of assessing water cycle modifications. The water stores are discussed in counterclockwise order in Figure 1a starting with atmospheric water while acknowledging that water stores are intimately and inherently interlinked, so considering them separately can be challenging. Evidence of local to regional regime shifts is ample and can potentially lead to nonlinear disruptions of the Earth system functions of water related to hydroclimatic and hydroecological regulation and storage through cross-scale interactions and cascading effects (Rocha et al., 2018; Steffen et al., 2018).

4.1. Atmospheric Water

The atmospheric water store fuels precipitation and is replenished by evapotranspiration and ocean evaporation. Here, we focus on the direct linkages between precipitation modification and hydroecological regime shifts, as well as evaporative change and regime shifts in hydroclimatic systems.

The Amazon rainforest is identified as a tipping element in the Earth system (Lenton et al., 2008; Steffen et al., 2018) with multiple alternative stable states primarily governed by precipitation (Hirota et al., 2011). A critical transition from evergreen rainforest to seasonal forest or savanna can have major consequences beyond the regional scale due to carbon release (Houghton et al., 2000), induced tipping of the South American monsoon system (Boers et al., 2017), precipitation reduction (Zemp et al., 2017), and biodiversity loss (Malhi et al., 2008). Climate change and deforestation critically undermine the resilience of the Amazon forest. The position of the climate change-induced threshold is uncertain due to a large spread in models' ability to simulate precipitation, fire feedback, and ecosystem response, among others (Cox et al., 2013; Huntingford et al., 2013; Nobre & Borma, 2009). The threshold of deforestation-induced Amazon forest dieback has been suggested to be between 10% and 40% depending on definitions and extent of forest transition considered (Nobre & Borma, 2009; Pires & Costa, 2013). The Congo rainforest and Southeast Asian rainforests are other less investigated tropical forest regions exhibiting similar regime shift mechanisms and consequences for Earth system functions as the Amazon forest (Bell et al., 2015; Lawrence & Vandecar, 2015; Staver et al., 2011). In temperate regions, drought conditions and considerable reductions in precipitation have been proven to trigger rapid coniferous forest declines in the southwestern United States (Anderegg et al., 2013). The tipping point has been found to be the persistence of an intense water

deficit over 11 months (Huang et al., 2015). Small changes in precipitation regimes are also known to have induced structural changes in wetland ecosystems, and abrupt ecological transitions in coastal wetlands are expected to expand to new coastal wetlands as hydroclimatic changes step up in the future (Osland et al., 2016).

Monsoons are large-scale seasonal reversals of atmospheric circulation with threshold behavior in terms of variability regime, atmospheric moisture transport, and spatial distribution (Lenton et al., 2008). Shifts in monsoon systems can have abrupt consequences at the continental scale (Lenton et al., 2008). For example, the West African monsoon shift had a major influence on the stable states between the Green Sahara regime (11,000–5,000 years ago) and the current Desert Sahara regime (Tierney et al., 2017; Yu et al., 2015). The rainy phase of monsoons brings large amounts of precipitation, turning landscapes green and replenishing rivers and aquifers, and is crucially important for agriculture and ecosystems. While large-scale tipping behavior of monsoon systems are primarily driven by global climate and ocean processes, regional land surface processes mediated by the water cycle are also one of several important factors that are able to influence monsoon dynamics. Studies have, for example, shown that increased evapotranspiration (i.e., latent heat) over irrigated land areas can decrease the land–ocean pressure gradient and thereby delay the Indian summer monsoon onset (e.g., Tuinenburg, 2013). In the Sahel, land degradation that reduces evapotranspiration and increases the north–south temperature gradient appears to enhance the African easterly jet, which drives moisture westward out of the region, in opposition to the southwesterly West African monsoon circulation (Hagos et al., 2014). Land feedback effects extend to tropics, and, for example, Nogherotto et al. (2013) showed that decreased evapotranspiration over deforested areas in the Congo has a seasonal influence on the strength of the west and south-equatorial African monsoon. Land–ocean gradient is also important in the South American monsoon system: Boers et al. (2017) showed that deforestation can induce a tipping point in the South American monsoon when latent heat release is no longer sufficient to maintain a positive feedback that enhances atmospheric inflow from the Atlantic, which would cause significant precipitation decline in the western Amazon and regions further downstream.

4.2. Soil Moisture

Soil moisture mediates terrestrial ecosystem transitions and desertification processes. Decrease in soil moisture caused by vegetation loss, erosion, and compaction creates a self-reinforcing feedback that prevents the re-establishment of plants (e.g., Karssenberg et al., 2017; Whitford et al., 2006) or causes shifts in ecosystem species composition (Loheide & Gorelick, 2007). Soil moisture-related land degradation has the potential for cascading and teleconnected impacts on the Earth's energy balance through, for example, large-scale albedo change and desert dust that follows wind beyond continents with effects on both climate systems and nutrient balance in distant regions (Bestelmeyer et al., 2015; Geist & Lambin, 2004). Also, deficits in soil moisture and changes in terrestrial water storage can severely diminish the primary production and carbon sequestration capacity of the terrestrial biosphere (Humphrey et al., 2018). Regions with important soil carbon storage and sequestration are the Northern Hemisphere that has the largest soil organic carbon stocks and the tropics that have seen the largest decrease in carbon stocks due to agricultural expansion, respectively (Cherlet et al., 2018).

4.3. Surface Water

While aquatic ecosystems can be negatively impacted by changes in streamflow (Carlisle et al., 2017; Gido et al., 2010; Perkin et al., 2017; Vörösmarty et al., 2010), there is no clear evidence or mechanism by which local- or basin-scale changes in aquatic biosphere integrity could scale up to have a planetary impact. However, one local-scale tipping point related to aquatic ecosystems is the transition of streams from perennial to intermittent, which can lead to a reorganization of local food webs (Bogan & Lytle, 2011). This transition is likely to be driven by changes in the groundwater storage function of water, which acts as a buffer against short-term hydroclimatic variability by providing a stable supply of baseflow to streams. A second local-scale hydroecological tipping point that has been identified in the literature is food web collapse associated with eutrophication and salinization. For the perennial–ephemeral and oligotrophic–eutrophic regime shifts, evidence of tipping points to eutrophic states (Wang & Temmerman, 2013) or even lake disappearance by water use-induced drying exist in several regions around the world, the most well known being the Aral Sea (Shibuo et al., 2007). We are not aware of studies that look beyond an individual body of water to trigger widespread shifts in Earth system function. Finally, various species- to ecosystem-

level effects, in particular for migratory species, have been documented in river ecosystems due to reductions in hydrologic connectivity (Pringle, 2003) caused by the global proliferation of anthropogenic dam and barrier construction (Grill et al., 2019).

Wetland ecosystems are rich in water-dependent biodiversity and play a multifaceted role for many Earth system processes, including high rates of evapotranspiration and groundwater recharge, temporary water storage, and sediment exchange. Large wetland complexes located downstream of streams and rivers may experience stress-induced tipping points due to variations in their hydrological characteristics. Also, coastal wetlands with mangrove ecosystems under such stress can experience reductions in their mangrove development and extensive mangrove mortality (Jimenez et al., 1985; Smith, 1992; Twilley & Rivera-Monroy, 2005); reductions of freshwater inputs to coastal wetlands or hydrological modification of their natural flows and connectivity due to reservoirs have already resulted in massive mangrove mortality episodes involving hypersalinity conditions in several wetlands around the world from which the wetlands have not been able to completely recover (Barreto, 2008; Cintron et al., 1978; Jaramillo et al., 2018; Jimenez et al., 1985).

Finally, surface water flows can affect Earth system processes due to their natural freshwater, sediment, and nutrient delivery to coastal zones and the ocean. Reductions in these flows, due to either climate or anthropogenic impacts including sediment trapping in reservoirs, may shift the balance between aggradation and erosion rates of large river deltas leading to land loss and cascading effects in marine ecosystems (Syvitski et al., 2009; Tessler et al., 2018), including their ability to sequester carbon (Duarte et al., 2004). Altered flows can potentially affect global ocean circulation systems through changes in salinity and temperature; for example, changes in Arctic runoff may affect Arctic ocean stratification, circulation, and ice cover (Nummelin et al., 2016) with implications for global oceanic circulation, including the Atlantic overturning meridional circulation and thermohaline circulation.

4.4. Groundwater

Several potential groundwater-related tipping points are associated with the storage function of groundwater. Most critical for aquatic ecosystems is the role of groundwater as a stable supply of baseflow, and therefore a key tipping point is when a stream transitions from perennial to intermittent (Bogan & Lytle, 2011) due to groundwater depletion (see section 4.3). However, groundwater-related tipping points are also present for terrestrial groundwater-dependent ecosystems. Groundwater within or near the root zone provides a stable supply of water, particularly during drought, for many natural and agricultural crops via capillary rise and direct groundwater uptake (Booth et al., 2016; Brown et al., 2011; Eamus et al., 2015; Rohde et al., 2017; Zipper et al., 2015, 2017). Numerous examples exist for critical transitions associated with regional-scale impacts of changes in groundwater storage, including groundwater depletion leading to riparian forest loss (Scott et al., 1999), rising groundwater levels leading to widespread flooding in Argentina (Houspanossian et al., 2016; Kuppel et al., 2015), and loss of dry forests leading to regional salinization in Australia (Clarke et al., 2002; George et al., 1999) and the Chaco region of Argentina (Giménez et al., 2016; Marchesini et al., 2017). Since groundwater is estimated to influence terrestrial ecosystems over 7–17% of global land area (Fan et al., 2013) and can contribute substantially to evapotranspiration (Lowry & Loheide, 2010; Soylyu et al., 2011, 2014; Yeh & Famiglietti, 2009), it likely constitutes an important component of terrestrial evapotranspiration. Thus, important groundwater-dependent ecosystems that may contribute to regional-scale shifts could potentially have a proportionally larger influence on Earth system dynamics. For instance, groundwater is an essential contributor to evapotranspiration in the Amazon basin (Fan et al., 2017; Miguez-Macho & Fan, 2012a, 2012b).

4.5. Frozen Water

Unlike the other water sub-boundaries, global tipping elements associated with frozen water storage have been studied extensively due to their potential contributions to global sea level rise (Intergovernmental Panel on Climate Change, 2019). While mass loss due to the melting of grounded glacial ice is widespread and accelerating in both the Antarctic and Greenland ice sheets (Velicogna et al., 2014), the West Antarctic Ice Sheet is thought to be vulnerable to tipping point-type dynamics, which would occur if ocean water was able to undercut the ice sheet and rapidly accelerate melt (Feldmann & Levermann, 2015; Lenton et al., 2008; Notz, 2009; Rignot et al., 2004). The collapse of the West Antarctic Ice Sheet would lead to an

estimated 5 m of sea level rise, which is comparable in magnitude with the total sea level change over the past ~7,000 years (Fleming et al., 1998). While the loss of Arctic sea ice would have impacts on regional and global climate due to reduced albedo and is a distinctive marker of alternate states of the Earth system, its melting sea ice would not impact sea levels (Bathiany et al., 2016; Notz, 2009; Tietsche et al., 2011). Widespread destabilization of permafrost is another potential tipping point related to frozen water (Lenton et al., 2008), as permafrost thaw leads to the release of greenhouse gases, which is a positive feedback on climate change and causes increasing sediment transport (Bring et al., 2016; Syvitski, 2002). There is increasing evidence for abrupt thaw mechanisms at local scales (Chasmer & Hopkinson, 2017; Chipman & Hu, 2017; Schuur et al., 2015; Zipper et al., 2018), though at global-scales permafrost thaw is thought to be a gradual source of carbon of approximately the same magnitude as land use change over the next century (Schuur et al., 2015).

5. Understanding Earth System Resilience Impacts of Global Water Cycle Modifications Using the Planetary Boundary Framework

5.1. Why Use the Planetary Boundary Framework?

Section 4 provides clues and hints of key regions and processes that could be important for identifying global water cycle modifications but does not deliver a systematic process for monitoring and detecting these. As discussed in section 2, the planetary boundary framework is, to our best knowledge, the only systematic effort for defining and monitoring the key processes and systems, which together regulate the state of the Earth system, the global water cycle being one of them. Monitoring human modifications of the global water cycle might be possible even without a systematic framework, but we argue that using a formalized framework, such as the planetary boundaries, is a pragmatic approach that will lead to scientific insight and implementation in governance and management while accounting for the intrinsic embedding of the water cycle in the Earth system. More significant scientific insights are possible because the planetary boundary framework rests on the foundations of decades of resilience research (described in section 2) and has been much discussed in multiple scientific communities. Additionally, this formalized framework links to other important current research areas in hydrology and other disciplines (described in section 1), which may lead to synergistic developments in multiple research strands. Stronger implementation in governance and management is also more likely as demonstrated by planetary boundaries already being considered in some governance and corporate management contexts (Clift et al., 2017; Galaz et al., 2012; Häyhä et al., 2016, 2018).

An additional benefit for governance and management is that the planetary boundary framework provides a useful bridge to water governance and management by formalizing the concept of the “safe operating space” for humanity. The safe operating space concept provides a set of quantitative scientific targets to keep the Earth within the relatively stable climatic conditions of the Holocene during which modern society developed. By setting the boundaries at a “safe” distance from scientifically defined dangerous levels or thresholds, the framework also involves normative judgements about how we choose to deal with risk and uncertainty. Various other water management indicators measure impact and status of water resources but do not explicitly connect these resources to Earth’s habitable conditions, such as water stress (Alcamo et al., 2007; Falkenmark, 1989; Smakhtin et al., 2004; Vorosmarty et al 2000), water depletion (Brauman et al., 2016), water scarcity (Brauman et al., 2016; Kummu et al., 2016), water footprints (Hoekstra & Mekonnen, 2012), groundwater footprints (Gleeson et al., 2012), water wedges (Wada et al., 2014), water use regimes (Weiskel et al., 2007), human appropriation of evapotranspiration (Gordon et al., 2005), and hydroclimatic separation (Destouni et al., 2012). The planetary boundaries framework complements this with information about the proximity of unwanted state shifts in Earth system stability, thus adding a simple aspirational metric to the toolbox.

Following Gleeson et al. (2019) we argue that although a planetary boundary for water is useful as a concept, the current definition and methodology does not adequately represent the role of water in the Earth system and should therefore be revised to reflect the key Earth system functions of water described in section 3. In the following subsections we propose an approach for a more robust analysis of global water cycle modifications using the planetary boundary framework, but similar approaches could be used to monitor and detect global water cycle modifications outside the framework.

5.2. Planetary Sub-Boundaries For Water Stores

In order to adequately represent the complexity and heterogeneity of the water cycle, we use the five stores (section 3) of water to divide the water cycle into planetary sub-boundaries that represent different processes and functions of water as proposed by Gleeson et al. (2019). Based on the five water stores, six planetary sub-boundaries (Figure 2) together represent the most important processes and crucial functions of water in the Earth system. Each store has a single sub-boundary based on a preliminary analysis of the core function of this store except atmospheric water that has two possible planetary sub-boundaries since atmospheric water has two distinct and important functions (hydroclimatic and hydroecologic).

The Earth system function and process addressed by each of the proposed sub-boundaries are shown in Figure 2 and summarized below:

- one atmospheric water sub-boundary focuses on the importance of evapotranspiration for climate pattern stability or land–atmosphere coupling stability (hydroclimatic regulation);
- second atmospheric water sub-boundary focuses on the role of precipitation in maintaining biomes, which is connected to biodiversity (hydroecologic regulation);
- soil moisture sub-boundary focuses on carbon uptake or net primary productivity (hydroclimatic regulation);
- surface water sub-boundary focuses on streamflow and related habitat that maintains aquatic biodiversity (hydroecologic regulation);
- groundwater sub-boundary focuses on baseflow that are important to aquatic biodiversity (hydroecologic regulation); and
- frozen water component focuses on ice sheet volume, which is important to sea level rise in the oceans (storage).


Possible control variables and suggested response variables are compiled in Figure 2. It is important to note that the proposed water planetary sub-boundaries do not represent the transport function of water, as these aspects are already considered in a separate “biogeochemical flows” planetary boundary. Outside the planetary boundaries framework, additional indicator(s) representing the transport function might need to be added.

5.3. Methodological Questions of Scale and Data


To monitor and detect global water cycle modifications, a number of methodological issues must be addressed. First are questions of space and time scales to consider in the analysis. Figure 2c summarizes the spatial aggregation appropriate for each of the suggested planetary sub-boundaries and respective control variables. For example, the surface water and groundwater components could be analysed at the large basin and regional aquifer scale (~10,000's to 100,000's km²), respectively. Time scales to consider depend on how the desirable baseline conditions against which current or future conditions may be compared are defined. The planetary boundary framework considers the Holocene epoch, yet robust global hydrologic data and models generally start in the ~1950s due to availability of widespread instrumental records and key data sets (Bierkens, 2015; Wada, 2016). This is also broadly consistent with the timing of the “great acceleration,” which is sometimes considered the onset of the post-Holocene Anthropocene (Steffen, Broadgate, et al., 2015; Zalasiewicz et al., 2015). We suggest ~1950s (or before if possible) as a Holocene-like “baseline” condition, understanding that this does not include all anthropogenic disturbances.

Second, a useful approach for monitoring and detecting global water cycle modifications will be model agnostic but does require some uniformity in the quantification approach. We argue that the appropriate approach requires explicit accounting of climate feedbacks, impacts on aquatic and terrestrial biodiversity, and other coupled impacts between the water cycle and Earth system stability in order to test relationships between control and response variables. Most existing global hydrologic models have only limited ability to simulate these feedbacks (Bierkens, 2015; Sood & Smakhtin, 2015). For example, global hydrological simulations used to quantify the current water planetary boundary (Steffen, Richardson, et al., 2015) were not dynamically coupled to a general circulation model. Recent study suggests that interactions among planetary boundaries predominantly aggravate human impacts on the Earth system and shrink the planetary “safe operating space” (Lade et al., 2019). It is likely that adequately assessing global water cycle

a) Core water functions for each water store



b) Dividing the water boundary into six sub-boundaries based on water functions



c) Key aspects of each of the proposed water sub-boundaries

	ATMOSPHERIC WATER (hydroclimate)	ATMOSPHERIC WATER (hydroecology)	SOIL MOISTURE (hydroclimate)	SURFACE WATER (hydroecology)	GROUNDWATER (hydroecology)	FROZEN WATER (storage)
Possible scale of analysis	Distributed hydrologic unit	Biomes or hydroclimatic regimes	Biomes or land cover groups	Large basins or river networks	Regional aquifers	Global
Possible response variable(s)	Climate pattern stability or land-atmosphere coupling stability	Terrestrial biosphere integrity (species richness or species/area)	Carbon uptake or net primary productivity	Aquatic biosphere integrity (species richness or species/area)	Terrestrial or aquatic biosphere integrity	Sea level rise
Possible interim planetary boundary	Percentage of global land area with evapotranspiration change within range of simulated future	Percentage of global land area with precipitation change within range of simulated future	Maintenance of global net primary productivity at or above levels under simulated future	Percentage of basins or total river length within environmental flow limits under simulated future	Percentage of basins with low flows meeting or exceeding simulated future	Volume of ice melt to keep sea level within limits under simulated future

Figure 2. (a) Each water store has core water function(s) to maintain Earth system resilience (see section 3), which could help quantify impacts of global water cycle modifications using (b) the planetary boundary framework. Defining water planetary sub-boundaries is based on the functional relationship between water stores and Earth system components. The green zone is the safe operating space; the yellow represents the zone of uncertainty; and the red indicates rising risks of large-scale, potentially abrupt and irreversible change (modified from Steffen, Richardson, et al., 2015). (c) Suggestions for key aspects of each of the six sub-boundaries including possible interim planetary boundary based on 2°C target for late this century in the United Nations Framework Convention on Climate Change Paris Agreement. The key water function for each sub-boundary is identified in parentheses (such as hydroecology for surface water).

modifications in the way we propose will necessitate revised models that robustly represent all water stores and their interactions with other parts of the Earth system or else better coupling with other models.

Input data in two different spatial perspectives may be useful for different water stores (see Weiskel et al., 2014 for longer discussion and definition): hydrologic units (distributed, open systems represented as pixels or raster cells) or semi-closed units (such as watersheds, aquifers, etc.). For example, for the surface water store, a semi-closed approach using large basins or river networks might be appropriate, whereas for the hydroclimatic function of the atmospheric water store (Figure 2), a distributed hydrologic system may be best since atmospheric water flows across traditional hydrological boundaries such as watershed divides

(e.g., Bosilovich, & Schubert, 2002; Dirmeyer et al., 2009; Van der Ent et al., 2011; Wang-Erlandsson et al., 2018, Chen et al., 2019). Also, for each water store, either hydrologic fluxes or the rate of change of hydrologic fluxes may be useful. The rate of change of hydrologic fluxes could be driven by a myriad of global change processes such as water use, land use, or climate change. An argument for the rate of change rather than the hydrologic fluxes is that the rate and time scale of environmental change may be more important to Earth system adaptation than absolute thresholds. Using streamflow as an example, aquatic ecosystems thrive in regions with a wide range of streamflow rates, but rapid change in streamflow could be problematic.

5.4. Spatial Analysis of Critical Regions of Global Water Cycle Modifications Using the Planetary Boundary Framework

To provide more methodology to water cycle modifications and water in the planetary boundary framework, we propose four new methods of spatial analysis that are based on the spatial distribution of impacts rather than summing of water fluxes (Figure 3). The current methodology for the freshwater use planetary boundary (or any other methodology that involves summing water fluxes) masks both the spatial and temporal heterogeneity of the water cycle and implies that resilience loss caused by water impacts in one place can be remediated by water abundance in another place. Each of these methods quantify the impact of water cycle modifications on Earth system resilience using control and response variables on the horizontal and vertical axis, respectively, and thus may also in the future be useful for setting water planetary sub-boundaries. Possible control and response variables for each water store are shown in Figure 2. The first three methods use input data described in section 5.3, whereas the last method uses previously mapped tipping elements (Lenton et al., 2008). The first three methods use four different types of data depending on the store: data in hydrologic units or semi-closed units, as well as data as fluxes or rate of change of fluxes (section 5.3).

1. The “unweighted approach” calculates the percentage of global land that has crossed a certain threshold using either the hydrologic fluxes or the hydrologic rate of change and the spatial perspective described in section 5.3. For example, for the surface water sub-boundary, the control variable could be the percentage of global land area of basins (or percentage length of river network to not bias by river length) not meeting environmental flow requirements; the input data would be at the scale of semi-closed units of basins or river networks (e.g., de Graaf et al., 2019). This approach would be useful if widespread degradation of conditions or change of fluxes or stores leads to significant change in the response variable. Another example is for the hydroclimatic function of atmospheric water store, where the percentage area that exceed a certain level of evapotranspiration change could be considered. A threshold could theoretically be set based on, for example, an evapotranspiration decrease, which induces an increase in surface temperature, reduction in atmospheric water store, and other impacts on local and regional climatic conditions. Here, we show in Figure 3a all evapotranspiration change that has occurred due to human induced land use change (e.g., including deforestation-induced decrease in evapotranspiration and irrigation-induced increase in evapotranspiration) as a possible basis for future threshold setting.
2. The “weighted approach” calculates the percentage of the global land area that has crossed a certain threshold weighted by the importance of that hydrologic unit to the Earth system function (also at a defined scale of analysis). This “weight” is multiplied with the data used in the “unweighted approach.” For example, again for the surface water sub-boundary, the control variable could be the percentage of global land area of basins not meeting environmental flow requirements weighted by aquatic biodiversity. This approach implies that there are regions where the Earth system function of water for the sub-boundary makes a more important contribution to the response variable. Another example is for the hydroclimatic function of atmospheric water store, where, for example, evapotranspiration change that has higher chance to influence downwind precipitation over land can be given a higher weight. Evapotranspiration changes at locations where a high percentage of evapotranspiration returns as precipitation over land also have higher changes to impact terrestrial rainfall, than places where the majority of evapotranspiration becomes oceanic precipitation. Thus, Figure 3b illustrates the evapotranspiration recycling ratio, the percentage of evapotranspiration that returns as precipitation over land, as a possible “weight.”
3. The “keystone region approach” identifies regions where certain water stores are disproportionately important to specific Earth system components. The concept of keystone regions is inspired by the concept of “keystone species,” a species that produces a major impact on their ecosystem and are considered




Figure 3. Alternative approaches for defining planetary boundary control and response variables (graphs on left hand side) using spatial analysis of critical regions of global water cycle modifications. Illustrative example using the hydroclimatic function of the atmospheric water store (maps on right hand side). The maps on the right side (a,b) are from Wang-Erlandsson et al., 2018 (based on modeled and reanalysis data over the period 2000–2013) and reproduced under Creative Commons Attribution 4.0 License. The map on the right side (c) is produced based directly from the (b) map. The map on the right side (d) map is from Lenton et al. (2008).

essential to maintaining optimum ecosystem function or structure (Mills et al., 1993), as well as the Pareto principle, also known as the 80–20 rule (Pareto, 1896). We hypothesize that a small number of regions (the “20” in the Pareto principle) have a disproportionate impact on the stability of the Earth system. We define a keystone region as a region where a water store produces a disproportionately important impact and could be essential to maintaining an Earth system component (e.g., atmospheric water in the Amazon is disproportionately important to the global climate system). “Disproportionately important” refers to the risk of direct or cascading impacts on other systems or regions following local or regional destabilization. For example, the Pareto principle could be used to identify the 20% of land area with the greatest weighting and exclude all other regions from global aggregation. For example, for the hydroclimatic function of atmospheric water store (Figure 3c), the areas with the highest evapotranspiration recycling ratio are considered keystone regions.

4. “Tipping elements” uses previously identified tipping elements in the Earth system (Lenton et al., 2008). Tipping elements are defined as subsystems of the Earth system that are at least subcontinental in scale and can be switched—under certain circumstances—into a qualitatively different state by small perturbations. The tipping point is the corresponding critical point—in forcing and a feature of the system—at which the future state of the system is qualitatively altered (Lenton et al., 2008) in a regime shift. For example, for the hydroclimatic function of atmospheric water store (Figure 3d), the monsoon systems could potentially be considered as atmospheric water tipping elements.

Different methods may be more effective or appropriate for each water store and water planetary sub-boundaries. A mixture of the most effective and appropriate methods for each water planetary sub-boundary could be used in setting the final planetary boundaries since the existing framework is based on a variety of different methods and metrics (Rockström et al., 2009a). Although the weighted approach, keystone region approach, and tipping element approach might all identify regions of hydrologic importance in global change, there are important differences between them. Tipping elements focus on cohesive subsystems and identifying regime shifts, whereas the keystone region approach focuses on cumulative and additive impacts across the Earth system, which may not be a cohesive subsystem and do not necessarily exhibit tipping behavior. The weighted approach also focuses on cumulative impacts but is less likely to result in cohesive regions than the keystone region method. All four approaches are different than other approaches in identifying regions of hydrologic importance in global change, like previous discussions of “water towers,” since the approaches here focus on changes to the Earth system, whereas “water towers” focus on the importance of specific regions for water resources.

5.5. Acting Upon Water Cycle Modifications by Setting and Using Water Planetary Sub-Boundaries

The process of setting fully elaborated planetary sub-boundaries with clearly defined relationships between control and response variables for the different water stores may take a considerable amount of time (at least ~5–10 years, comparable to other global change science synthesis activities). Yet there is significant interest in using the water planetary boundary, so we explored setting interim planetary sub-boundaries based on global standards for carbon and existing global data. Interim planetary boundaries for water could be set by quantifying the change in proposed control variables for each water component under the representative concentration pathways with related emissions and land use scenarios consistent with the United Nations Framework Convention on Climate Change Paris Agreement. In other words, these are the water boundaries that would arise if global carbon governance actors considered water impacts. These new boundaries could support water governance that increasingly addresses global issues through global water initiatives across sectors (Varady et al., 2009). Theoretical exploration of global water governance highlights a combination of multilevel design with a strong global dimension (Hoekstra, 2006; Pahl-Wostl et al., 2008). Global water governance could also be an integral part of a proposed Earth system governance framework (Biermann et al., 2012), integrated into existing global carbon governance or ideally developed as another parallel form of global governance.

We suggest that the water planetary boundary provides an effective framework that can integrate with and complement existing water management approaches at subglobal scales such as watersheds, aquifers, or nations. Subglobal use of the water planetary boundary is the focus of separate work (Zipper et al., 2020), and we only briefly introduce it here to highlight the potential utility of the water planetary boundary. Previous work with the planetary boundary framework at subglobal scales has either attempted to

calculate a “fair share” of the planetary boundary value allocated to a subglobal domain, for example, a nation (Häyhä et al., 2018) or use the control–response variable relationship foundational to the planetary boundary framework to develop local boundaries in a conceptually consistent manner (e.g., Dearing et al., 2014). Most critically, the planetary boundary framework allows water managers to account for potential global Earth system impacts of local water cycle modifications (e.g., local responsibility for global environmental challenges), a perspective not captured in existing water management frameworks, and provides a systematic framework for intercomparison across watershed, countries, companies, or other subglobal units where water is managed (Zipper et al., 2020). Given the lack of global water management and governance organizations, effective integration of water management across local- to global-scales may require innovative governance structures and approaches (Biermann, 2012; Galaz et al., 2012).

6. Concluding With an Invitation to Meet a Grand Challenge

The core functions of hydroclimatic regulation, hydroecological regulation, storage, and transport illuminate how water stores (atmospheric water, soil moisture, surface water, groundwater, and frozen water) are inextricably interconnected with Earth system components such as the atmosphere, land, and ocean through processes, mechanisms, and variables that are familiar to all hydrologists such as evapotranspiration, albedo, ice melt, streamflow, and so forth. The scientific and ethical grand challenge of examining water cycle modifications for Earth system resilience are motivated by the numerous research questions we introduced in section 1. The grand challenge first focuses on which water-related changes may lead to supraregional or global tipping points or gradual adverse transitions in critical water and Earth system functions (sections 2 and 3). A related question inherently relates to scaling between regional and global processes for which we provide new methods of analysis (section 5.4). Setting new water planetary sub-boundaries (section 5.5) may enable managing trade-offs between global development (e.g., the sustainable development goals) and increasing pressure on global water resources, which may motivate the development of water governance mechanisms and institutions that respond to and influence global water cycling. Finally, interactions and feedbacks from other systems such as food and energy drive many of these processes, so focusing on water cycle modifications may provide insights or tools for other initiatives.

We invite the hydrology and water resource community to apply serious and sustained attention toward understanding water cycle modifications and Earth system resilience, which could be transformative to our understanding of sociohydrologic systems across scales, up to the global, and provide a new approach for global hydrology modeling and analysis. We suggest three initiatives that can be tackled immediately and simultaneously by highly collaborative working groups from diverse backgrounds:

- Initiative 1 could compare the “weighted approach” to the “keystone region approach,” which could uncover differences in regions that are disproportionately important to different Earth system functions of water.
- Initiative 2, focusing on the rate of change of fluxes, could uncover the regions of the world experiencing the most rapid rates of change and investigate whether these have meaningful impact on different Earth system functions of water.
- Initiative 3 could identify and provisionally quantify interim, spatially explicit planetary sub-boundaries (which may not be possible or robust for all the planetary sub-boundaries).

Together, these three initiatives would lay the foundation for developing fully elaborated water planetary sub-boundaries and illuminating water cycle modifications in the Anthropocene. This ambitious scientific agenda also directly leads to important water policy implications. We therefore end with an invitation to the hydrology and water resources community to join us in following this grand challenge, which would initiate numerous interesting scientific journeys and help set precautionary planetary boundaries for water that reflect its undeniable importance in global sustainability and Earth system science.

Acknowledgments

This community project was supported by a STINT internationalization grant, Swedish Research Council (VR, project 2015-06503) and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning FORMAS (942-2015-740). We thank many members of the community who contributed to the discussions. We have no conflicts of interest and no data are used in this manuscript.

References

- Abraham, J. P., Baringer, M., Bindoff, N. L., Boyer, T., Cheng, L. J., Church, J. A., et al. (2013). A review of global ocean temperature observations: Implications for ocean heat content estimates and climate change. *Reviews of Geophysics*, *51*(3), 450–483. <https://doi.org/10.1002/rog.20022>
- Acreman, M. C., Overton, I. C., King, J., Wood, P. J., Cowx, I. G., Dunbar, M. J., et al. (2014). The changing role of ecohydrological science in guiding environmental flows. *Hydrological Sciences Journal*, *59*(3–4), 433–450. <https://doi.org/10.1080/02626667.2014.886019>

- Aeschbach-Hertig, W., & Gleeson, T. (2012). Regional strategies for the accelerating global problem of groundwater depletion. *Nature Geoscience*, 5(12), 853–861. <https://doi.org/10.1038/ngeo1617>
- Alcamo, J., Flörke, M., & Märker, M. (2007). Future long-term changes in global water resources driven by socio-economic and climatic changes. *Hydrological Sciences Journal*, 52(2), 247–275.
- Alcamo, J. M., Vörösmarty, C. J., Naiman, R. J., Lettenmaier, D. P., & Pahl-Wostl, C. (2008). A grand challenge for freshwater research: Understanding the global water system. *Environmental Research Letters*, 3(1), 010202.
- Anderegg, W. R. L., Kane, J. M., & Anderegg, L. D. L. (2013). Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change*, 3(1), 30–36. <https://doi.org/10.1038/nclimate1635>
- Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., et al. (2018). 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nature Communications*, 9(1), 1–12. <https://doi.org/10.1038/s41467-017-02771-y>
- Archfield, S. A., Clark, M., Arheimer, B., Hay, L. E., McMillan, H., Kiang, J. E., et al. (2015). Accelerating advances in continental domain hydrologic modeling. *Water Resources Research*, 51(12), 10,078–10,091. <https://doi.org/10.1002/2015WR017498>
- Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O., Swartz, B., Quental, T. B., et al. (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, 471(7336), 51–57.
- Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., et al. (2012). Approaching a state shift in Earth's biosphere. *Nature*, 486(7401), 52–58. <https://doi.org/10.1038/nature11018>
- Barona, E., Ramankutty, N., Hyman, G., & Coomes, O. T. (2010). The role of pasture and soybean in deforestation of the Brazilian Amazon. *Environmental Research Letters*, 5(2), 024002. <https://doi.org/10.1088/1748-9326/5/2/024002>
- Barreto, M. B. (2008). Diagnostics about the state of mangroves in Venezuela: Case studies from the National Park Morrocoy and Wildlife Refuge Cuare. In P. H. Lieth, D. M. G. Sucre, & B. Herzog (Eds.), *Mangroves and Halophytes: Restoration and Utilisation*, (pp. 51–64). Netherlands: Springer. https://doi.org/10.1007/978-1-4020-6720-4_6
- Bathiany, S., Notz, D., Mauritsen, T., Raedel, G., & Brovkin, V. (2016). On the potential for abrupt Arctic winter sea ice loss. *Journal of Climate*, 29(7), 2703–2719. <https://doi.org/10.1175/JCLI-D-15-0466.1>
- Bell, J. P., Tompkins, A. M., Bouka-Biona, C., & Sanda, I. S. (2015). A process-based investigation into the impact of the Congo basin deforestation on surface climate. *Journal of Geophysical Research*, 120(12), 5721–5739. <https://doi.org/10.1002/2014JD022586>
- Belmont, P., & Foufoula-Georgiou, E. (2017). Solving water quality problems in agricultural landscapes: New approaches for these non-linear, multiprocess, multiscale systems. *Water Resources Research*, 53(4), 2585–2590. <https://doi.org/10.1002/2017WR020839>
- Bestelmeyer, B. T., Williamson, J. C., Archer, S. R., Sayre, N. F., Duniway, M. C., Okin, G. S., & Herrick, J. E. (2015). Desertification, land use, and the transformation of global drylands. *Frontiers in Ecology and the Environment*, 13(1), 28–36. <https://doi.org/10.1890/140162>
- Bhaduri, A., Bogardi, J., Siddiqi, A., Voigt, H., Vörösmarty, C., Pahl-Wostl, C., et al. (2016). Achieving sustainable development goals from a water perspective. *Frontiers in Environmental Science*, 4, 64.
- Biddanda, B. A. (2017). Global significance of the changing freshwater carbon cycle. *Eos*, 98. <https://doi.org/10.1029/2017EO069751>
- Bierkens, M. F., & Wada, Y. (2019). Non-renewable groundwater use and groundwater depletion: A review. *Environmental Research Letters*, 14(6), 063002.
- Bierkens, M. F. P. (2015). Global hydrology 2015: State, trends, and directions. *Water Resources Research*, 51(7), 4923–4947. <https://doi.org/10.1002/2015WR017173>
- Bierkens, M. F. P., Bell, V. A., Burek, P., Chaney, N., Condon, L. E., David, C. H., et al. (2015). Hyper-resolution global hydrological modelling: What is next? *Hydrological Processes*, 29(2), 310–320. <https://doi.org/10.1002/hyp.10391>
- Biermann, F., Abbott, K., Andresen, S., Bäckstrand, K., Bernstein, S., Betsill, M. M., et al. (2012). Navigating the Anthropocene: Improving Earth system governance. *Science*, 335(6074), 1306–1307. <https://doi.org/10.1126/science.1217255>
- Biggs, R., Peterson, G., & Rocha, J. (2018). The regime shifts database: A framework for analyzing regime shifts in social-ecological systems. *Ecology and Society*, 23(3). <https://doi.org/10.5751/ES-10264-230309>
- Black, P. E. (1997). Watershed functions. *JAWRA Journal of the American Water Resources Association*, 33(1), 1–11.
- Boers, N., Marwan, N., Barbosa, H. M. J., & Kurths, J. (2017). A deforestation-induced tipping point for the South American monsoon system. *Scientific Reports*, 7, 41489. <https://doi.org/10.1038/srep41489>
- Bogan, M. T., & Lytle, D. A. (2011). Severe drought drives novel community trajectories in desert stream pools. *Freshwater Biology*, 56(10), 2070–2081. <https://doi.org/10.1111/j.1365-2427.2011.02638.x>
- Booth, E. G., & Loheide, S. P. (2012). Comparing surface effective saturation and depth-to-water-level as predictors of plant composition in a restored riparian wetland. *Ecohydrology*, 5(5), 637–647. <https://doi.org/10.1002/eco.250>
- Booth, E. G., Zipper, S. C., Loheide, S. P., & Kucharik, C. J. (2016). Is groundwater recharge always serving us well? Water supply provisioning, crop production, and flood attenuation in conflict in Wisconsin, USA. *Ecosystem Services*, 21, Part A, 153–165. <https://doi.org/10.1016/j.ecoser.2016.08.007>
- Bosilovich, M. G., & Schubert, S. D. (2002). Water vapor tracers as diagnostics of the regional hydrologic cycle. *Journal of Hydrometeorology*, 3(2), 149–165.
- Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. *Elem Sci Anth*, 4.
- Bring, A., Fedorova, I., Dibike, Y., Hinzman, L., Mard, J., Mernild, S. H., et al. (2016). Arctic terrestrial hydrology: A synthesis of processes, regional effects, and research challenges. *Journal of Geophysical Research – Biogeosciences*, 121(3), 621–649. <https://doi.org/10.1002/2015JG003131>
- Brown, C. M., Lund, J. R., Cai, X., Reed, P. M., Zagana, E. A., Ostfeld, A., et al. (2015). The future of water resources systems analysis: Toward a scientific framework for sustainable water management. *Water Resources Research*, 51(8), 6110–6124. <https://doi.org/10.1002/2015WR017114>
- Brown, J., Bach, L., Aldous, A., Wyers, A., & DeGagné, J. (2011). Groundwater-dependent ecosystems in Oregon: An assessment of their distribution and associated threats. *Frontiers in Ecology and the Environment*, 9(2), 97–102. <https://doi.org/10.1890/090108>
- Cai, X., Wallington, K., Shafiee-Jood, M., & Marston, L. (2018). Understanding and managing the food-energy-water nexus—Opportunities for water resources research. *Advances in Water Resources*, 111, 259–273. <https://doi.org/10.1016/j.advwatres.2017.11.014>
- Carlisle, D. M., Grantham, T. E., Eng, K., & Wolock, D. M. (2017). Biological relevance of streamflow metrics: Regional and national perspectives. *Freshwater Science*, 36(4), 927–940. <https://doi.org/10.1086/694913>
- Catford, J. A., Morris, W. K., Vesk, P. A., Gippel, C. J., & Downes, B. J. (2014). Species and environmental characteristics point to flow regulation and drought as drivers of riparian plant invasion. *Diversity and Distributions*, 20(9), 1084–1096. <https://doi.org/10.1111/ddi.12225>

- Chase-Dunn, C., and T. D. Hall. 1997. Ecological degradation and the evolution of world-systems. *Journal of World-Systems Research* 3(3):403-431. Available online at: <http://jwsr.ucr.edu/archive/vol3/v3n3a3.php>.
- Chasmer, L., & Hopkinson, C. (2017). Threshold loss of discontinuous permafrost and landscape evolution. *Global Change Biology*, 23(7), 2672–2686. <https://doi.org/10.1111/gcb.13537>
- Chen, X., Leung, L. R., Wigmosta, M., & Richmond, M. (2019). Impact of atmospheric rivers on surface hydrological processes in western US watersheds. *Journal of Geophysical Research: Atmospheres*.
- Cherlet, M., Hutchinson, C., Reynolds, J., Hill, J., Sommer, S., & von Maltitz, G. (Eds) (2018). *World Atlas of desertification*. Luxemburg: Publication Office of the European Union.
- Chipman, M. L., & Hu, F. S. (2017). Linkages among climate, fire, and thermoerosion in Alaskan tundra over the past three millennia. *Journal of Geophysical Research – Biogeosciences*, 122(12), 3362–3377. <https://doi.org/10.1002/2017JG004027>
- Cintron, G., Lugo, A. E., Pool, D. J., & Morris, G. (1978). Mangroves of arid environments in Puerto Rico and adjacent islands. *Biotropica*, 10(2), 110–121. <https://doi.org/10.2307/2388013>
- Clarke, C. J., George, R. J., Bell, R. W., & Hatton, T. J. (2002). Dryland salinity in south-western Australia: Its origins, remedies, and future research directions. *Soil Research*, 40(1), 93–113.
- Clift, R., Sim, S., King, H., Chenoweth, J. L., Christie, I., Clavreul, J., et al. (2017). The challenges of applying planetary boundaries as a basis for strategic decision-making in companies with global supply chains. *Sustainability*, 9(2), 279. <https://doi.org/10.3390/su9020279>
- Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., & Luke, C. M. (2013). Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. *Nature*, 494(7437), 341–344. <https://doi.org/10.1038/nature11882>
- Dalin, C., Wada, Y., Kastner, T., & Puma, M. J. (2017). Groundwater depletion embedded in international food trade. *Nature*, 543(7647), 700–704. <https://doi.org/10.1038/nature21403>
- D'Almeida, C., Vörösmarty, C. J., Hurr, G. C., Marengo, J. A., Dingman, S. L., & Keim, B. D. (2007). The effects of deforestation on the hydrological cycle in Amazonia: A review on scale and resolution. *International Journal of Climatology*, 27(5), 633–647. <https://doi.org/10.1002/joc.1475>
- de Graaf, I. E. M., Gleeson, T., van Beek, L. P. H. (Rens), Sutanudjaja, E. H., & Bierkens, M. F. P. (2019). Environmental flow limits to global groundwater pumping. *Nature*, 574(7776), 90–94. <https://doi.org/10.1038/s41586-019-1594-4>
- Deane, D. C., Nicol, J. M., Gehrig, S. L., Harding, C., Aldridge, K. T., Goodman, A. M., & Brookes, J. D. (2017). Hydrological-niche models predict water plant functional group distributions in diverse wetland types. *Ecological Applications*, 27(4), 1351–1364. <https://doi.org/10.1002/eap.1529>
- Dearing, J. A., Wang, R., Zhang, K., Dyke, J. G., Haberl, H., Hossain, M. S., et al. (2014). Safe and just operating spaces for regional social-ecological systems. *Global Environmental Change*, 28, 227–238. <https://doi.org/10.1016/j.gloenvcha.2014.06.012>
- Denemark, R. A., Friedman, J., Gills, B. K., & Modelski, G. (Eds) (2000). *World system history: The social science of long-term change*. London, UK: Routledge.
- Destouni, G., Jaramillo, F., & Prieto, C. (2012). Hydroclimatic shifts driven by human water use for food and energy production. *Nature Climate Change*, 3(3), 213–217. <https://doi.org/10.1038/nclimate1719>
- Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J. L., & Blöschl, G. (2013). Socio-hydrology: Conceptualising human-flood interactions. *Hydrology and Earth System Sciences*, 17(8), 3295–3303. <https://doi.org/10.5194/hess-17-3295-2013>
- Dickerson-Lange, S. E., & Mitchell, R. (2014). Modeling the effects of climate change projections on streamflow in the Nooksack River basin, Northwest Washington. *Hydrological Processes*, 28(20), 5236–5250. <https://doi.org/10.1002/hyp.10012>
- Dingman, S. L. (2002). *Physical hydrology*. Upper Saddle River: Prentice-Hall.
- Dirmeyer, P. A., Schlosser, C. A., & Brubaker, K. L. (2009). Precipitation, recycling, and land memory: An integrated analysis. *Journal of Hydrometeorology*, 10(1), 278–288.
- Donges, J. F., & Barfuss, W. (2017). From math to metaphors and back again: Social-ecological resilience from a multi-agent-environment perspective. *GAIA - Ecological Perspectives for Science and Society*, 26(1), 182–190. <https://doi.org/10.14512/gaia.26.S1.5>
- Duarte, C. M., Middelburg, J. J., & Caraco, N. (2004). Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences Discussions*, 1(1), 659–679.
- Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015). Projections of future meteorological drought and wet periods in the Amazon. *Proceedings of the National Academy of Sciences*, 112(43), 13172–13177. <https://doi.org/10.1073/pnas.1421010112>
- Eamus, D., Zolfaghar, S., Villalobos-Vega, R., Cleverly, J., & Huete, A. (2015). Groundwater-dependent ecosystems: Recent insights from satellite and field-based studies. *Hydrology and Earth System Sciences*, 19(10), 4229–4256. <https://doi.org/10.5194/hess-19-4229-2015>
- Earle, A., Cascao, A. E., Hansson, S., Jägerskog, A., Swain, A., Öjendal, J., Öjendal, J. (2015). Transboundary water management and the climate change debate. Routledge. <https://doi.org/10.4324/9780203098929>
- Ellis, S., Mellor, T., & Mellor, T. (2002). *Soils and environment*. Routledge. <https://doi.org/10.4324/9780203415245>
- Falkenmark, M. (1989). The massive water scarcity now threatening Africa: Why isn't it being addressed? *Ambio*, 112–118.
- Falkenmark, M., Wang-Erlandsson, L., & Rockström, J. (2019). Understanding of water resilience in the Anthropocene. *Journal of Hydrology X*, 2, 100009.
- Fan, Y., Clark, M., Lawrence, D. M., Swenson, S., Band, L. E., Brantley, S. L., et al. (2019). Hillslope hydrology in global change research and earth system modeling. *Water Resources Research*, 55(2), 1737–1772. <https://doi.org/10.1029/2018WR023903>
- Fan, Y., Leung, L. R., Duan, Z., Wigmosta, M. S., Maxwell, R. M., Chambers, J. Q., & Tomasella, J. (2017). Influence of landscape heterogeneity on water available to tropical forests in an Amazonian catchment and implications for modeling drought response. *Journal of Geophysical Research-Atmospheres*, 122(16), 2017JD027066. <https://doi.org/10.1002/2017JD027066>
- Fan, Y., Li, H., & Miguez-Macho, G. (2013). Global patterns of groundwater table depth. *Science*, 339(6122), 940–943. <https://doi.org/10.1126/science.1229881>
- Feldmann, J., & Levermann, A. (2015). Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin. *Proceedings of the National Academy of Sciences*, 112(46), 14,191–14,196. <https://doi.org/10.1073/pnas.1512482112>
- Fleming, K., Johnston, P., Zwart, D., Yokoyama, Y., Lambeck, K., & Chappell, J. (1998). Refining the eustatic sea-level curve since the Last Glacial maximum using far- and intermediate-field sites. *Earth and Planetary Science Letters*, 163(1), 327–342. [https://doi.org/10.1016/S0012-821X\(98\)00198-8](https://doi.org/10.1016/S0012-821X(98)00198-8)
- Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., & Holling, C. S. (2004). Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution, and Systematics*, 35(1), 557–581. <https://doi.org/10.1146/annurev.ecolsys.35.021103.105711>
- Galaz, V., Biermann, F., Folke, C., Nilsson, M., & Olsson, P. (2012). Global environmental governance and planetary boundaries: An introduction. *Ecological Economics*, 81, 1–3. <https://doi.org/10.1016/j.ecolecon.2012.02.023>

- Galaz, V., Crona, B., Dauriach, A., Scholtens, B., & Steffen, W. (2018). Finance and the Earth system – Exploring the links between financial actors and non-linear changes in the climate system. *Global Environmental Change*, 53, 296–302. <https://doi.org/10.1016/j.gloenvcha.2018.09.008>
- Geist, H. J., & Lambin, E. F. (2004). Dynamic causal patterns of desertification. *Bioscience*, 54(9), 817–829. [https://doi.org/10.1641/0006-3568\(2004\)054\[0817:DCPOD\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0817:DCPOD]2.0.CO;2)
- George, R. J., Nulsen, R. A., Ferdowsian, R., & Raper, G. P. (1999). Interactions between trees and groundwaters in recharge and discharge areas - A survey of Western Australian sites. *Agricultural Water Management*, 39(2–3), 91–113. [https://doi.org/10.1016/S0378-3774\(98\)00073-0](https://doi.org/10.1016/S0378-3774(98)00073-0)
- Gerten, D. (2013). A vital link: Water and vegetation in the Anthropocene. *Hydrology and Earth System Sciences*, 17(10), 3841–3852. <https://doi.org/10.5194/hess-17-3841-2013>
- Gerten, D., Hoff, H., Rockström, J., Jägermeyr, J., Kummu, M., & Pastor, A. V. (2013). Towards a revised planetary boundary for consumptive freshwater use: role of environmental flow requirements. *Current Opinion in Environmental Sustainability*, 5(6), 551–558. <https://doi.org/10.1016/j.cosust.2013.11.001>
- Gido, K. B., Dodds, W. K., & Eberle, M. E. (2010). Retrospective analysis of fish community change during a half-century of landuse and streamflow changes. *Journal of the North American Benthological Society*, 29(3), 970–987.
- Giménez, R., Mercáu, J., Nosoletto, M., Páez, R., & Jobbágy, E. (2016). The ecohydrological imprint of deforestation in the semiarid Chaco: Insights from the last forest remnants of a highly cultivated landscape. *Hydrological Processes*, 30(15), 2603–2616. <https://doi.org/10.1002/hyp.10901>
- Gleeson, T., Erlandsson, L. W., Zipper, S., Porkka, M., Jaramillo F., Gerten D., et al. (2019). The water planetary boundary: Interrogation and revision. Preprint published at <https://eartharxiv.org/swhma/>
- Gleeson, T., & Richter, B. (2018). How much groundwater can we pump and protect environmental flows through time? Presumptive standards for conjunctive management of aquifers and rivers. *River Research and Applications*, 34(1), 83–92. <https://doi.org/10.1002/rra.3185>
- Gleeson, T., Wada, Y., Bierkens, M. F. P., & van Beek, L. P. H. (2012). Sustainable water balance of global aquifers revealed by groundwater footprint. *Nature*, 488, 197–200. <https://doi.org/10.1038/nature11295>
- Gleick, P. H. (2000). *The world's water 2000-2001*. Washington: Island Press.
- Gordon, L. J., Steffen, W., Jönsson, B. F., Folke, C., Falkenmark, M., & Johannessen, Å. (2005). Human modification of global water vapor flows from the land surface. *Proceedings of the National Academy of Sciences of the United States of America*, 102(21), 7612–7617. <https://doi.org/10.1073/pnas.0500208102>
- Gotts, N. M. (2007). Resilience, panarchy, and world-systems analysis. *Ecology and Society*, 12(1). <https://doi.org/24>
- Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., et al. (2019). Mapping the world's free-flowing rivers. *Nature*, 569(7755), 215–221. <https://doi.org/10.1038/s41586-019-1111-9>
- Guilod, B. P., Orlovsky, B., Miralles, D. G., Teuling, A. J., & Seneviratne, S. I. (2015). Reconciling spatial and temporal soil moisture effects on afternoon rainfall. *Nature Communications*, 6, 6443. <https://doi.org/10.1038/ncomms7443>
- Hall, T. D. (Ed) (2000). *A world-systems reader: New perspectives on gender, urbanism, cultures, indigenous peoples and ecology*. London, UK: Rowman and Littlefield.
- Häyhä, T., Cornell, S. E., Hoff, H., Lucas, P., & van Vuuren, D. (2018). *Operationalizing the concept of a safe operating space at the EU level - First steps and explorations (Stockholm Resilience Centre Technical Report, prepared in collaboration with Stockholm Environment Institute (SEI) and PBL Netherlands Environmental Assessment Agency)*. Stockholm Resilience Centre: Stockholm University, Sweden.
- Häyhä, T., Lucas, P. L., van Vuuren, D. P., Cornell, S. E., & Hoff, H. (2016). From planetary boundaries to national fair shares of the global safe operating space — How can the scales be bridged? *Global Environmental Change*, 40, 60–72. <https://doi.org/10.1016/j.gloenvcha.2016.06.008>
- Hecky, R. E., Mugidde, R., Ramlal, P. S., Talbot, M. R., & Kling, G. W. (2010). Multiple stressors cause rapid ecosystem change in Lake Victoria. *Freshwater Biology*, 55(s1), 19–42. <https://doi.org/10.1111/j.1365-2427.2009.02374.x>
- Heistermann, M. (2017). HESS Opinions: A planetary boundary on freshwater use is misleading. *Hydrology and Earth System Sciences*, 21(7), 3455–3461. <https://doi.org/10.5194/hess-21-3455-2017>
- Henshaw, P. C., Charlson, R. J., & Burges, S. J. (2000). 6 - Water and the Hydrosphere. In M. C. Jacobson, R. J. Charlson, H. Rodhe, & G. H. Orians (Eds.), *International Geophysics* (Vol. 72, pp. 109–131). London, UK: Elsevier Academic Press. [https://doi.org/10.1016/S0074-6142\(00\)80112-6](https://doi.org/10.1016/S0074-6142(00)80112-6)
- Henszey, R. J., Pfeiffer, K., & Keough, J. R. (2004). Linking surface-and ground-water levels to riparian grassland species along the Platte River in Central Nebraska, USA. *Wetlands*, 24(3), 665–687.
- Hirota, M., Holmgren, M., Van Nes, E. H., & Scheffer, M. (2011). Global resilience of tropical forest and savanna to critical transitions. *Science*, 334(6053), 232–235.
- Hoek, W. Z. (2008). The last glacial-interglacial transition. *Episodes*, 31(2), 226–229.
- Hoekstra, A. Y. (2006). The global dimension of water governance: Nine reasons for global arrangements in order to cope with local water problems. Retrieved from <https://research.utwente.nl/en/publications/the-global-dimension-of-water-governance-nine-reasons-for-global->
- Hoekstra, A. Y., & Mekonnen, M. M. (2012). The water footprint of humanity. *Proceedings of the National Academy of Sciences*, 109(9), 3232–3237. <https://doi.org/10.1073/pnas.1109936109>
- Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, 4(1), 1–23. <https://doi.org/10.1146/annurev.es.04.110173.000245>
- Holling, C. S., & Gunderson, L. H. (2002). Resilience and adaptive cycles. In L. H. Gunderson, & C. S. Holling (Eds.), *Panarchy: Understanding transformations in human and natural systems*, (pp. 25–62). Washington, D.C., USA: Island Press.
- Houghton, R. A., Skole, D. L., Nobre, C. A., Hackler, J. L., Lawrence, K. T., & Chomentowski, W. H. (2000). Annual fluxes of carbon from deforestation and regrowth in the Brazilian Amazon. *Nature*, 403(6767), 301.
- Houspanossian, J., Kuppel, S., Nosoletto, M., Bella, C. D., Oricchio, P., Barrucand, M., et al. (2016). Long-lasting floods buffer the thermal regime of the Pampas. *Theoretical and Applied Climatology*, 131(1-2), 111–120. <https://doi.org/10.1007/s00704-016-1959-7>
- Huang, K., Yi, C., Wu, D., Zhou, T., Zhao, X., Blanford, W. J., et al. (2015). Tipping point of a conifer forest ecosystem under severe drought. *Environmental Research Letters*, 10(2), 024011. <https://doi.org/10.1088/1748-9326/10/2/024011>
- Humphrey, V., Zscheischler, J., Ciais, P., Gudmundsson, L., Sitch, S., & Seneviratne, S. I. (2018). Sensitivity of atmospheric CO₂ growth rate to observed changes in terrestrial water storage. *Nature*, 560(7720), 628. <https://doi.org/10.1038/s41586-018-0424-4>

- Huntingford, C., Zelazowski, P., Galbraith, D., Mercado, L. M., Sitch, S., Fisher, R., et al. (2013). Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nature Geoscience*, 6(4), 268–273. <https://doi.org/10.1038/ngeo1741>
- Huntington, T. G. (2006). Evidence for intensification of the global water cycle: Review and synthesis. *Journal of Hydrology*, 319(1), 83–95. <https://doi.org/10.1016/j.jhydrol.2005.07.003>
- Huntington, T. G., Weiskel, P. K., Wolock, D. M., & McCabe, G. J. (2018). A new indicator framework for quantifying the intensity of the terrestrial water cycle. *Journal of Hydrology*, 559, 361–372. <https://doi.org/10.1016/j.jhydrol.2018.02.048>
- Immerzeel, W. W., van Beek, L. P. H., & Bierkens, M. F. P. (2010). Climate change will affect the Asian water towers. *Science*, 328(5984), 1382–1385. <https://doi.org/10.1126/science.1183188>
- IPCC (2013). Climate change 2013: The physical science basis.
- IPCC (2019). Special report on the ocean and cryosphere in a changing climate.
- Jaramillo, F., & Destouni, G. (2015a). Comment on “Planetary boundaries: Guiding human development on a changing planet.”. *Science*, 348(6240), 1217–1217. <https://doi.org/10.1126/science.aaa9629>
- Jaramillo, F., & Destouni, G. (2015b). Local flow regulation and irrigation raise global human water consumption and footprint. *Science*, 350(6265), 1248–1251. <https://doi.org/10.1126/science.aad1010>
- Jaramillo, F., Licero, L., Åhlen, I., Manzoni, S., Rodríguez-Rodríguez, J. A., Guittard, A., et al. (2018). Effects of hydroclimatic change and rehabilitation activities on salinity and mangrove recovery in the Ciénaga Grande de Santa Marta. *Colombia. Wetlands*, 1–13. <https://doi.org/10.1007/s13157-018-1024-7>
- Jimenez, J. A., Lugo, A. E., & Cintron, G. (1985). Tree mortality in mangrove forests. *Biotropica*, 177–185.
- Karsenberg, D., Bierkens, M. F. P., & Rietkerk, M. (2017). Catastrophic shifts in semiarid vegetation-soil systems may unfold rapidly or slowly. *The American Naturalist*, 190(6), E145–E155. <https://doi.org/10.1086/694413>
- Kummu, M., Guillaume, J. H. A., De Moel, H., Eisner, S., Flörke, M., Porkka, M., et al. (2016). The world’s road to water scarcity: Shortage and stress in the 20th century and pathways towards sustainability. *Scientific Reports*, 6(1), 38495. <https://doi.org/10.1038/srep38495>
- Kuppel, S., Houspanossian, J., Noretto, M. D., & Jobbágy, E. G. (2015). What does it take to flood the Pampas?: Lessons from a decade of strong hydrological fluctuations. *Water Resources Research*, 51(4), 2937–2950. <https://doi.org/10.1002/2015WR016966>
- Lade, S. J., Steffen, W., de Vries, W., Carpenter, S. R., Donges, J. F., Gerten, D., et al. (2019). Human impacts on planetary boundaries amplified by Earth system interactions. *Nature Sustainability*, 1–10.
- Lawrence, D., & Vandecar, K. (2015). Effects of tropical deforestation on climate and agriculture. *Nature Climate Change*, 5(1), 27–36. <https://doi.org/10.1038/nclimate2430>
- Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., & Schellnhuber, H. J. (2008). Tipping elements in the Earth’s climate system. *Proceedings of the National Academy of Sciences*, 105(6), 1786–1793. <https://doi.org/10.1073/pnas.0705414105>
- Loheide, S. P., & Gorelick, S. M. (2007). Riparian hydroecology: A coupled model of the observed interactions between groundwater flow and meadow vegetation patterning. *Water Resources Research*, 43(7).
- Lowry, C. S., & Loheide, S. P. (2010). Groundwater-dependent vegetation: Quantifying the groundwater subsidy. *Water Resources Research*, 46(6), W06202. <https://doi.org/10.1029/2009WR008874>
- Lund, J. R. (2015). Integrating social and physical sciences in water management. *Water Resources Research*, 51(8), 5905–5918. <https://doi.org/10.1002/2015WR017125>
- Malhi, Y., Roberts, J. T., Betts, R. A., Killeen, T. J., Li, W., & Nobre, C. A. (2008). Climate change, deforestation, and the fate of the Amazon. *Science*, 319(5860), 169–172.
- Marchesini, V. A., Giménez, R., Noretto, M. D., & Jobbágy, E. G. (2017). Ecohydrological transformation in the dry Chaco and the risk of dryland salinity: Following Australia’s footsteps? *Ecohydrology*, 10(4), e1822. <https://doi.org/10.1002/eco.1822>
- Marshall, C. R. (2006). Explaining the Cambrian “explosion” of animals. *Annual Review of Earth and Planetary Sciences*, 34, 355–384.
- McGuire, K. J., & McDonnell, J. J. (2006). A review and evaluation of catchment transit time modeling. *Journal of Hydrology*, 330(3–4), 543–563.
- Medema, W., McIntosh, B., & Jeffrey, P. (2008). From premise to practice: A critical assessment of integrated water resources management and adaptive management approaches in the water sector. *Ecology and Society*, 13(2), art29. <https://doi.org/10.5751/ES-02611-130229>
- Meyfroidt, P., Rudel, T. K., & Lambin, E. F. (2010). Forest transitions, trade, and the global displacement of land use. *Proceedings of the National Academy of Sciences*, 107(49), 20917–20922.
- Micklin, P. (2007). The Aral Sea disaster. *Annual Review of Earth and Planetary Sciences*, 35(1), 47–72. <https://doi.org/10.1146/annurev.earth.35.031306.140120>
- Micklin, P., Aladin, N. V., & Plotnikov, I. (2016). Aral Sea. Springer. <https://www.springer.com/gp/book/9783642023552>
- Miguez-Macho, G., & Fan, Y. (2012a). The role of groundwater in the Amazon water cycle: 1. Influence on seasonal streamflow, flooding and wetlands. *Journal of Geophysical Research-Atmospheres*, 117(D15), n/a. <https://doi.org/10.1029/2012JD017539>
- Miguez-Macho, G., & Fan, Y. (2012b). The role of groundwater in the Amazon water cycle: 2. Influence on seasonal soil moisture and evapotranspiration. *Journal of Geophysical Research-Atmospheres*, 117(D15), D15114. <https://doi.org/10.1029/2012JD017540>
- Mills, L. S., Soulé, M. E., & Doak, D. F. (1993). The keystone-species concept in ecology and conservation. *Bioscience*, 43(4), 219–224.
- Mitchell, J. F. B. (1989). The “greenhouse” effect and climate change. *Reviews of Geophysics*, 27(1), 115–139. <https://doi.org/10.1029/RG027i001p00115>
- Montanari, A., Young, G., Savenije, H. H. G., Hughes, D., Wagener, T., Ren, L. L., et al. (2013). “Panta Rhei—Everything Flows”: Change in hydrology and society—The IAHS Scientific Decade 2013–2022. *Hydrological Sciences Journal*, 58(6), 1256–1275. <https://doi.org/10.1080/02626667.2013.809088>
- Motew, M., Chen, X., Booth, E. G., Carpenter, S. R., Pinkas, P., Zipper, S. C., et al. (2017). The Influence of Legacy P on lake water quality in a Midwestern Agricultural Watershed. *Ecosystems*, 20(8), 1468–1482. <https://doi.org/10.1007/s10021-017-0125-0>
- National Marine Fisheries Service (2016). *Coastal multispecies recovery plan*. Santa Rosa, CA: National Marine Fisheries Service, West Coast Region.
- National Research Council (1991). *Opportunities in the hydrologic sciences*. Washington, DC: The National Academies Press. Retrieved from <https://doi.org/10.17226/1543>
- Nepstad, D. C., de Carvalho, C. R., Davidson, E. A., Jipp, P. H., Lefebvre, P. A., Negreiros, G. H., et al. (1994). The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures. *Nature*, 372(6507), 666–669. <https://doi.org/10.1038/372666a0>
- Nobre, A. D. (2014). *The future climate of Amazonia*. INPA and ARA, São José Dos Campos Brazil: Scientific Assessment Report. Sponsored by CCST-INPE.

- Nobre, C. A., & Borma, L. D. S. (2009). 'Tipping points' for the Amazon forest. *Current Opinion in Environmental Sustainability*, 1(1), 28–36. <https://doi.org/10.1016/j.cosust.2009.07.003>
- Nogherotto, R., Coppola, E., Giorgi, F., & Mariotti, L. (2013). Impact of Congo Basin deforestation on the African monsoon. *Atmospheric Science Letters*, 14(1), 45–51. <https://doi.org/10.1002/asl2.416>
- Noss, R. F., Dobson, A. P., Baldwin, R., Beier, P., Davis, C. R., Dellasala, D. A., et al. (2012). Bolder thinking for conservation. *Conservation Biology*, 26(1), 1–4. <https://doi.org/10.1111/j.1523-1739.2011.01738.x>
- Notz, D. (2009). The future of ice sheets and sea ice: Between reversible retreat and unstoppable loss. Proceedings of the National Academy of Sciences, pnas.0902356106. <https://doi.org/10.1073/pnas.0902356106>
- Nummelin, A., Ilicak, M., Li, C., & Smedsrud, L. H. (2016). Consequences of future increased Arctic runoff on Arctic Ocean stratification, circulation, and sea ice cover. *Journal of Geophysical Research, Oceans*, 121(1), 617–637.
- Oki, T., & Kanae, S. (2006). Global hydrological cycles and world water resources. *Science*, 313(5790), 1068–1072. <https://doi.org/10.1126/science.1128845>
- Oki, T., Yano, S., & Hanasaki, N. (2017). Economic aspects of virtual water trade. *Environmental Research Letters*, 12(4), 044002. <https://doi.org/10.1088/1748-9326/aa625f>
- Osland, M. J., Enwright, N. M., Day, R. H., Gabler, C. A., Stagg, C. L., & Grace, J. B. (2016). Beyond just sea-level rise: Considering macroclimatic drivers within coastal wetland vulnerability assessments to climate change. *Global Change Biology*, 22(1), 1–11. <https://doi.org/10.1111/gcb.13084>
- Overton, I. C., Jolly, I. D., Slavich, P. G., Lewis, M. M., & Walker, G. R. (2006). Modelling vegetation health from the interaction of saline groundwater and flooding on the Chowilla floodplain, South Australia. *Australian Journal of Botany*, 54(2), 207–220. <https://doi.org/10.1071/BT05020>
- Pahl-Wostl, C., Gupta, J., & Petry, D. (2008). Governance and the global water system: A theoretical exploration. *Global Governance: A Review of Multilateralism and International Organizations*, 14(4), 419–435.
- Pardini, R., de Arruda Bueno, A., Gardner, T. A., Prado, P. I., & Metzger, J. P. (2010). Beyond the fragmentation threshold hypothesis: Regime shifts in biodiversity across fragmented landscapes. *PLoS ONE*, 5(10), e13666. <https://doi.org/10.1371/journal.pone.0013666>
- Pareto, V. (1896). Course of political economy. Lausanne.
- Perkin, J. S., Gido, K. B., Falke, J. A., Fausch, K. D., Crockett, H., Johnson, E. R., & Sanderson, J. (2017). Groundwater declines are linked to changes in Great Plains stream fish assemblages. Proceedings of the National Academy of Sciences, 201618936. Retrieved from <http://www.pnas.org/content/early/2017/06/20/1618936114.short>
- Pires, G. F., & Costa, M. H. (2013). Deforestation causes different subregional effects on the Amazon bioclimatic equilibrium. *Geophysical Research Letters*, 40(14), 3618–3623. <https://doi.org/10.1002/grl.50570>
- Poff, N. L., & Matthews, J. H. (2013). Environmental flows in the Anthropocene: Past progress and future prospects. *Current Opinion in Environmental Sustainability*, 5(6), 667–675.
- Poff, N. L., Richter, B. D., Arthington, A. H., Bunn, S. E., Naiman, R. J., Kendy, E., et al. (2009). The ecological limits of hydrologic alteration (ELOHA): A new framework for developing regional environmental flow standards. *Freshwater Biology*, 55(1), 147–170. Retrieved from <https://doi.org/10.1111/j.1365-2427.2009.02204.x>
- Pool, T. K., Olden, J. D., Whittier, J. B., & Paukert, C. P. (2010). Environmental drivers of fish functional diversity and composition in the Lower Colorado River Basin. *Canadian Journal of Fisheries and Aquatic Sciences*, 67(11), 1791–1807. <https://doi.org/10.1139/F10-095>
- Porkka, M., Kumm, M., Siebert, S., & Flörke, M. (2012). The role of virtual water flows in physical water scarcity: The case of Central Asia. *International Journal of Water Resources Development*, 28(3), 453–474. <https://doi.org/10.1080/07900627.2012.684310>
- Pringle, C. (2003). What is hydrologic connectivity and why is it ecologically important? *Hydrological Processes*, 17(13), 2685–2689. <https://doi.org/10.1002/hyp.5145>
- Reager, J. T., Gardner, A. S., Famiglietti, J. S., Wiese, D. N., Eicker, A., & Lo, M.-H. (2016). A decade of sea level rise slowed by climate-driven hydrology. *Science*, 351(6274), 699–703. <https://doi.org/10.1126/science.aad8386>
- Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A., & Thomas, R. (2004). Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf. *Geophysical Research Letters*, 31(18). <https://doi.org/10.1029/2004GL020697>
- Rocha, J. C., Peterson, G., Bodin, Ö., & Levin, S. (2018). Cascading regime shifts within and across scales. *Science*, 362(6421), 1379–1383.
- Rocha, J. C., Peterson, G. D., & Biggs, R. (2015). Regime shifts in the Anthropocene: Drivers, risks, and resilience. *PLoS ONE*, 10(8), e0134639. <https://doi.org/10.1371/journal.pone.0134639>
- Rockström, J., Falkenmark, M., Folke, C., Lannerstad, M., Barron, J., Enfors, E., et al. (2014). Water resilience for human prosperity. Cambridge University Press.
- Rockström, J., Gordon, L. J., Folke, C., Falkenmark, M., & Engwall, M. (1999). Linkages among water vapor flows, food production, and terrestrial ecosystem services. *Conservation Ecology*, 3(2), 5.
- Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S., Lambin, E. F., et al. (2009b). A safe operating space for humanity. *Nature*, 461(7263), 472–475. <https://doi.org/10.1038/461472a>
- Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S. I., Lambin, E., et al. (2009a). Planetary boundaries: Exploring the safe operating space for humanity. *Ecology and Society*, 14(2), art32. <https://doi.org/10.5751/ES-03180-140232>
- Rodhe, H. (1990). A comparison of the contribution of various gases to the greenhouse effect. *Science*, 248(4960), 1217–1219. <https://doi.org/10.1126/science.248.4960.1217>
- Rohde, M. M., Freund, R., & Howard, J. (2017). A Global synthesis of managing groundwater dependent ecosystems under sustainable groundwater policy. *Groundwater*, n/a-n/a. <https://doi.org/10.1111/gwat.12511>
- Rullí, M. C., Savioli, A., & D'Odorico, P. (2013). Global land and water grabbing. *Proceedings of the National Academy of Sciences*, 110(3), 892–897.
- Savenije, H. H., Hoekstra, A. Y., & van der Zaag, P. (2014). Evolving water science in the Anthropocene. *Hydrology and Earth System Sciences*, 18(1), 319–332.
- Scheffer, M., Carpenter, S., Foley, J. A., Folke, C., & Walker, B. (2001). Catastrophic shifts in ecosystems. *Nature*, 413(6856), 591–596. <https://doi.org/10.1038/35098000>
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., et al. (2015). Climate change and the permafrost carbon feedback. *Nature*, 520(7546), 171–179. <https://doi.org/10.1038/nature14338>
- Scott, M. L., Shafroth, P. B., & Auble, G. T. (1999). Responses of riparian cottonwoods to alluvial water table declines. *Environmental Management*, 23(3), 347–358. <https://doi.org/10.1007/s002679900191>

- Shibuo, Y., Jarsjö, J., & Destouni, G. (2007). Hydrological responses to climate change and irrigation in the Aral Sea drainage basin. *Geophysical Research Letters*, *34*(21). <https://doi.org/10.1029/2007GL031465>
- Shiklomanov, I. A., & Rodda, J. C. (2003). World water resources at the beginning of the twenty-first century. Retrieved from <http://bases.bireme.br/cgi-bin/wxislind.exe/iah/online/?IsisScript=iah/iah.xis&src=google&base=REPIDISCA&lang=p&nextAction=lnk&exprSearch=30315&indexSearch=ID>
- Sivapalan, M., Konar, M., Srinivasan, V., Chhatre, A., Wutich, A., Scott, C. A., et al. (2014). Socio-hydrology: Use-inspired water sustainability science for the Anthropocene. *Earth's Future*, *2*(4), 225–230. <https://doi.org/10.1002/2013EF000164>
- Sivapalan, M., Savenije, H. H., & Blöschl, G. (2012). Socio-hydrology: A new science of people and water. *Hydrological Processes*, *26*(8), 1270–1276.
- Smakhtin, V. U., Revenga, C., & Döll, P. (2004). A pilot global assessment of environmental water requirements and scarcity. *Water International*, *29*(3), 307–317.
- Smith, T. J. (1992). Forest Structure. In A. I. Robertson & D. M. Alongi (Eds.), *Tropical mangrove ecosystems* (pp. 101–136). Washington DC: American Geophysical Union. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1029/CE041p0101/summary>
- Smith, V. H., & Schindler, D. W. (2009). Eutrophication science: Where do we go from here? *Trends in Ecology & Evolution*, *24*(4), 201–207.
- Sood, A., & Smakhtin, V. (2015). Global hydrological models: A review. *Hydrological Sciences Journal*, *60*(4), 549–565.
- Soylu, M. E., Istanbuloglu, E., Lenters, J. D., & Wang, T. (2011). Quantifying the impact of groundwater depth on evapotranspiration in a semi-arid grassland region. *Hydrology and Earth System Sciences*, *15*(3), 787–806. <https://doi.org/10.5194/hess-15-787-2011>
- Soylu, M. E., Kucharik, C. J., & Loheide, S. P. (2014). Influence of groundwater on plant water use and productivity: Development of an integrated ecosystem – Variably saturated soil water flow model. *Agricultural and Forest Meteorology*, *189–190*, 198–210. <https://doi.org/10.1016/j.agrformet.2014.01.019>
- Spera, S. A., Galford, G. L., Coe, M. T., Macedo, M. N., & Mustard, J. F. (2016). Land-use change affects water recycling in Brazil's last agricultural frontier. *Global Change Biology*, *22*(10), 3405–3413. <https://doi.org/10.1111/gcb.13298>
- Staal, A., Tuinenburg, O. B., Bosmans, J. H. C., Holmgren, M., van Nes, E. H., Scheffer, M., et al. (2018). Forest-rainfall cascades buffer against drought across the Amazon. *Nature Clim Change*, *8*, 539–543. <https://doi.org/10.1038/s41558-018-0177-y>
- Staver, A. C., Archibald, S., & Levin, S. A. (2011). The global extent and determinants of savanna and forest as alternative biome states. *Science*, *334*(6053), 230–232. <https://doi.org/10.1126/science.1210465>
- Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O., & Ludwig, C. (2015). The trajectory of the Anthropocene: The great acceleration. *The Anthropocene Review*, *2*(1), 81–98. <https://doi.org/10.1177/2053019614564785>
- Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., et al. (2015). Planetary boundaries: Guiding human development on a changing planet. *Science*, *347*(6223), 1259855. <https://doi.org/10.1126/science.1259855>
- Steffen, W., Rockström, J., Richardson, K., Lenton, T. M., Folke, C., Liverman, D., et al. (2018). Trajectories of the earth system in the Anthropocene. *Proceedings of the National Academy of Sciences*, *115*(33), 8252–8259. <https://doi.org/10.1073/pnas.1810141115>
- Sterling, S. M., Ducharme, A., & Polcher, J. (2012). The impact of global land-cover change on the terrestrial water cycle. *Nature Climate Change*, *3*(4), 385–390. <https://doi.org/10.1038/nclimate1690>
- Summerfield, M. A. (2005). The changing landscape of geomorphology. *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group*, *30*(6), 779–781.
- Summerfield, M. A. (2014). *Global geomorphology*. New York, USA: Routledge.
- Swann, A. L. S., Longo, M., Knox, R. G., Lee, E., & Moorcroft, P. R. (2015). Future deforestation in the Amazon and consequences for South American climate. *Agricultural and Forest Meteorology*, *214–215*, 12–24. <https://doi.org/10.1016/j.agrformet.2015.07.006>
- Syvitski, J. P., Kettner, A. J., Overeem, I., Hutton, E. W., Hannon, M. T., Brakenridge, G. R., et al. (2009). Sinking deltas due to human activities. *Nature Geoscience*, *2*(10), 681–686. <https://doi.org/10.1038/ngeo629>
- Syvitski, J. P. M. (2002). Sediment discharge variability in Arctic rivers: Implications for a warmer future. *Polar Research*, *21*(2), 323–330. <https://doi.org/10.1111/j.1751-8369.2002.tb00087.x>
- Qiuhong Tang, & Taikan Oki (Eds.). (2016). Terrestrial water cycle and climate change: Natural and human-induced impacts, In *Geophysical Monograph 221*, First Edition. (pp. 3–16). American Geophysical Union.
- Tessler, Z. D., Vörösmarty, C. J., Overeem, I., & Syvitski, J. P. (2018). A model of water and sediment balance as determinants of relative sea level rise in contemporary and future deltas. *Geomorphology*, *305*, 209–220.
- Tierney, J. E., Pausata, F. S. R., & De Menocal, P. B. (2017). Rainfall regimes of the green sahara. *Science Advances*, *3*(1). <https://doi.org/10.1126/sciadv.1601503>
- Tietsche, S., Notz, D., Jungclaus, J. H., & Marotzke, J. (2011). Recovery mechanisms of Arctic summer sea ice. *Geophysical Research Letters*, *38*(2). <https://doi.org/10.1029/2010GL045698>
- Tölggyessy, J. (1993). Chemistry and biology of water, air and soil: Environmental aspects (Vol. 53). Elsevier.
- Tuinenburg, O. A. (2013). Atmospheric effects of irrigation in moonsoon climate: The Indian subcontinent (PhD Thesis). Wageningen University.
- Twilley, R. R., & Rivera-Monroy, V. H. (2005). Developing performance measures of mangrove wetlands using simulation models of hydrology, nutrient biogeochemistry, and community dynamics. *Journal of Coastal Research*, 79–93. Retrieved from <http://www.jstor.org/stable/25736617>
- UNESCO (1978). *World water balance and water resources of the earth (Vol. USSR committee for the international hydrologic decade)*. Paris: UNESCO.
- van der Ent, R. J., Savenije, H. G. G., Schaeffli, B., & Steele-Dunne, S. C. (2010). Origin and fate of atmospheric moisture over continents. *Water Resources Research*, *46*(9), 1–12. <https://doi.org/10.1029/2010WR009127>
- Van der Ent, R. J., & Savenije, H. H. G. (2011). Length and time scales of atmospheric moisture recycling. *Atmospheric Chemistry and Physics*, *11*(5), 1853–1863.
- Varady, R. G., Meehan, K., & McGovern, E. (2009). Charting the emergence of 'global water initiatives' in world water governance. *Physics and Chemistry of the Earth, Parts A/B/C*, *34*(3), 150–155. <https://doi.org/10.1016/j.pce.2008.06.004>
- Velicogna, I., Sutterley, T. C., & van den Broeke, M. R. (2014). Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data. *Geophysical Research Letters*, *41*(22), 8130–8137. <https://doi.org/10.1002/2014GL061052>
- Vilà-Guerau de Arellan, J., van Heerwaarden, C. C., van Stratum, B. J. H., & van den Dries, K. (2015). *Atmospheric boundary layer integrating air chemistry and land interactions: Atmospheric science and meteorology*. New York, USA: Cambridge University Press.
- Vogel, R. M., Lall, U., Cai, X., Rajagopalan, B., Weiskel, P. K., Hooper, R. P., & Matalas, N. C. (2015). Hydrology: The interdisciplinary science of water. *Water Resources Research*, *51*(6), 4409–4430. <https://doi.org/10.1002/2015WR017049>

- Vörösmarty, C. J., Hoekstra, A. Y., Bunn, S. E., Conway, D., & Gupta, J. (2015). Fresh water goes global. *Science*, *349*(6247), 478–479. <https://doi.org/10.1126/science.aac6009>
- Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., et al. (2010). Global threats to human water security and river biodiversity. *Nature*, *467*(7315), 555–561. <https://doi.org/10.1038/nature09440>
- Wada, Y. (2016). Modeling groundwater depletion at regional and global scales: Present state and future prospects. *Surveys in Geophysics*, *37*(2), 419–451. <https://doi.org/10.1007/s10712-015-9347-x>
- Wada, Y., Gleeson, T., & Esnault, L. (2014). Wedge approach to water stress. *Nature Geoscience*, *7*(9), 615–617. <https://doi.org/10.1038/ngeo2241>
- Wada, Y., Lo, M.-H., Yeh, P. J.-F., Reager, J. T., Famiglietti, J. S., Wu, R.-J., & Tseng, Y.-H. (2016). Fate of water pumped from underground and contributions to sea-level rise. *Nature Climate Change*, *6*(8), 777–780. <https://doi.org/10.1038/nclimate3001>
- Wada, Y., van Beek, L. P., Sperna Weiland, F. C., Chao, B. F., Wu, Y.-H., & Bierkens, M. F. (2012). Past and future contribution of global groundwater depletion to sea-level rise. *Geophysical Research Letters*, *39*(9).
- Wagner, T., Sivapalan, M., Troch, P., & Woods, R. (2007). Catchment classification and hydrologic similarity. *Geography Compass*, *1*(4), 901–931. <https://doi.org/10.1111/j.1749-8198.2007.00039.x>
- Wagner, T., Sivapalan, M., Troch, P. A., McGlynn, B. L., Harman, C. J., Gupta, H. V., et al. (2010). The future of hydrology: An evolving science for a changing world. *Water Resources Research*, *46*(5). <https://doi.org/10.1029/2009WR008906>
- Walker, B., Holling, C. S., Carpenter, S., & Kinzig, A. (2004). Resilience, adaptability and transformability in social–ecological systems. *Ecology and Society*, *9*(2).
- Wallerstein, I. M. (1974). The modern world-system. Volume 1. In *Capitalist agriculture and the origins of the European world-economy in the sixteenth century* (pp. 2–355). Cambridge, UK: Cambridge University Press.
- Wang, C., & Temmerman, S. (2013). Does biogeomorphic feedback lead to abrupt shifts between alternative landscape states?: An empirical study on intertidal flats and marshes, 118, 229–240. <https://doi.org/10.1029/2012JF002474>
- Wang-Erlandsson, L., Fetzer, I., Keys, P. W., van der Ent, R. J., Savenije, H. H. G., & Gordon, L. J. (2018). Remote land use impacts on river flows through atmospheric teleconnections. *Hydrology and Earth System Sciences*, *22*(8), 4311–4328. <https://doi.org/10.5194/hess-22-4311-2018>
- Watson, K. A., Mayer, A. S., & Reeves, H. W. (2014). Groundwater availability as constrained by hydrogeology and environmental flows. *Groundwater*, *52*(2), 225–238. <https://doi.org/10.1111/gwat.12050>
- Weiskel, P. K., Vogel, R. M., Steeves, P. A., Zarriello, P. J., DeSimone, L. A., & Ries, K. G., III. (2007). Water use regimes: Characterizing direct human interaction with hydrologic systems. *Water Resources Research*, *43*(4), W04402. Retrieved from <https://doi.org/10.1029/2006WR005062>
- Whitford, W. G., Virginia, R. A., Huenneke, L. F., Reynolds, J. F., Schlesinger, W. H., Jarrell, W. M., & Cunningham, G. L. (2006). Biological feedbacks in global desertification. *Science*, *247*(4946), 1043–1048. <https://doi.org/10.1126/science.247.4946.1043>
- Wiens, J. A., Moss, M. R., Fahrig, L., & Milne, B. (2005). *Issues and perspectives in landscape ecology*. Cambridge, UK: Cambridge University Press.
- Wood, E. F., Roundy, J. K., Troy, T. J., van Beek, L. P. H., Bierkens, M. F. P., Blyth, E., et al. (2011). Hyperresolution global land surface modeling: Meeting a grand challenge for monitoring Earth's terrestrial water. *Water Resources Research*, *47*(5). <https://doi.org/10.1029/2010WR010090>
- Xu, J., Grumbine, R. E., Shrestha, A., Eriksson, M., Yang, X., Wang, Y., & Wilkes, A. (2009). The melting Himalayas: Cascading effects of climate change on water, biodiversity, and livelihoods. *Conservation Biology*, *23*(3), 520–530. <https://doi.org/10.1111/j.1523-1739.2009.01237.x>
- Yang, Y., Saatchi, S. S., Xu, L., Yu, Y., Choi, S., Phillips, N., et al. (2018). Post-drought decline of the Amazon carbon sink. *Nature Communications*, *9*(1), 1, 3172–9. <https://doi.org/10.1038/s41467-018-05668-6>
- Yeh, P. J.-F., & Famiglietti, J. S. (2009). Regional groundwater evapotranspiration in Illinois. *Journal of Hydrometeorology*, *10*(2), 464–478. <https://doi.org/10.1175/2008JHM1018.1>
- Yu, K., D'Odorico, P., Bhattachan, A., Okin, G. S., & Evan, A. T. (2015). Dust-rainfall feedback in West African Sahel. *Geophysical Research Letters*, *42*(18), 7563–7571. <https://doi.org/10.1002/2015GL065533>
- Zalasiewicz, J., Waters, C. N., Williams, M., Barnosky, A. D., Cearreta, A., Crutzen, P., et al. (2015). When did the Anthropocene begin? A mid-twentieth century boundary level is stratigraphically optimal. *Quaternary International*, *383*, 196–203. <https://doi.org/10.1016/j.quaint.2014.11.045>
- Zemp, D. C., Schleussner, C.-F., Barbosa, H. M. J., Hirota, M., Montade, V., Sampaio, G., et al. (2017). Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks. *Nature Communications*, *8*(1), 14681. <https://doi.org/10.1038/ncomms14681>
- Zipper, S. C., Jaramillo, F., Erlandsson, L. W., Cornell, S., Gleeson, T., Porkka, M., et al. (2020). Integrating the water planetary boundary with water management from local to global scales. *Earth's Future*, e2019EF001377. <https://doi.org/10.1029/2019EF001377>
- Zipper, S. C., Lamontagne-Hallé, P., McKenzie, J. M., & Rocha, A. V. (2018). Groundwater controls on post-fire permafrost thaw: Water and energy balance effects. *Journal of Geophysical Research - Earth Surface*, *123*(10), 2677–2694. <https://doi.org/10.1029/2018JF004611>
- Zipper, S. C., Soylu, M. E., Booth, E. G., & Loheide, S. P. (2015). Untangling the effects of shallow groundwater and soil texture as drivers of subfield-scale yield variability. *Water Resources Research*, *51*(8), 6338–6358. <https://doi.org/10.1002/2015WR017522>
- Zipper, S. C., Soylu, M. E., Kucharik, C. J., & Loheide, S. P. II (2017). Quantifying indirect groundwater-mediated effects of urbanization on agroecosystem productivity using MODFLOW-AgroIBIS (MAGI), a complete critical zone model. *Ecological Modelling*, *359*, 201–219. <https://doi.org/10.1016/j.ecolmodel.2017.06.002>