

ACCEPTED MANUSCRIPT • OPEN ACCESS

Peatland protection and restoration are key for climate change mitigation

To cite this article before publication: Florian Humpenöder *et al* 2020 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/abae2a>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2020 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Peatland protection and restoration are key for climate change mitigation

Florian Humpenöder^{1*}, Kristine Karstens^{1,2}, Hermann Lotze-Campen^{1,2}, Jens Leifeld³, Lorenzo Menichetti⁴, Alexandra Barthelmes⁵, Alexander Popp¹

¹Potsdam Institute for Climate Impact Research, Germany.

²Humboldt University of Berlin, Germany.

³Agroscope, Switzerland.

⁴Swedish University of Agricultural Sciences, Sweden.

⁵Greifswald Mire Centre, Germany.

*Corresponding author

humpenoeder@pik-potsdam.de

Accepted Manuscript

Abstract

Peatlands cover only about 3% the global land area, but store about twice as much carbon as global forest biomass. If intact peatlands are drained for agriculture or other human uses, peat oxidation can result in considerable CO₂ emissions and other greenhouse gases (GHG) for decades or even centuries. Despite their importance, emissions from degraded peatlands have so far not been included explicitly in mitigation pathways compatible with the Paris Agreement. Such pathways include land-demanding mitigation options like bioenergy or afforestation with substantial consequences for the land system. Therefore, besides GHG emissions owing to the historic conversion of intact peatlands, the increased demand for land in current mitigation pathways could result in drainage of presently intact peatlands, e.g. for bioenergy production. Here, we present the first quantitative model-based projections of future peatland dynamics and associated GHG emissions in the context of a 2°C mitigation pathway. Our spatially explicit land-use modelling approach with global coverage simultaneously accounts for future food demand, based on population and income projections, and land-based mitigation measures. Without dedicated peatland policy and even in the case of peatland protection, our results indicate that the land system would remain a net source of CO₂ throughout the 21st century. This result is in contrast to the outcome of current mitigation pathways, in which the land system turns into a net carbon sink by 2100. However, our results indicate that it is possible to reconcile land use and GHG emissions in mitigation pathways through a peatland protection *and* restoration policy. According to our results, the land system would turn into a global net carbon sink by 2100, as projected by current mitigation pathways, if about 60%

1
2
3 of present-day degraded peatlands would be rewetted in the coming decades, next to the
4
5 protection of intact peatlands.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript

Introduction

Peatlands cover only about 3% the global land area (381-463 Mha) but store about twice as much carbon as global forest biomass (Leifeld and Menichetti 2018, Leifeld *et al* 2019, Yu *et al* 2010, Joosten 2010, Joosten *et al* 2016). About 43-51 Mha of the global peatland area is degraded (Leifeld and Menichetti 2018, Leifeld *et al* 2019, Joosten 2010), i.e. those peatlands have been drained for agriculture, forestry or peat extraction. Degraded peatlands emit carbon dioxide (CO₂) from peat oxidation and other greenhouse gases (GHG) for centuries if not rewetted (Leifeld and Menichetti 2018, Wilson *et al* 2016). Estimates of present-day global GHG emissions from degraded peatlands range between 1.30-1.91 Gt CO_{2eq} yr⁻¹ (Leifeld and Menichetti 2018, Leifeld *et al* 2019, Joosten 2010), which corresponds to 2.6-3.8% of total global anthropogenic GHG emissions (49 Gt CO_{2eq} yr⁻¹ in 2010) (Edenhofer *et al* 2014).

Natural peatlands accumulate peat from dead plant material, thereby removing CO₂ from the atmosphere (Hooijer *et al* 2010). At the same time, the decomposition of dead plant material under anaerobic conditions causes methane (CH₄) emissions with a GHG effect roughly 30 times that of CO₂ (Jauhiainen *et al* 2016). Over a 100-year timescale these counteracting effects can even cancel out each other (Barthelmes *et al* 2015). Therefore, natural peatlands can be considered from a moderate net GHG sink (Levy and Gray 2015) to climate neutral (Barthelmes *et al* 2015). When peatlands are drained the peat starts to decompose aerobically and therefore at a much higher rate compared to the anaerobic accumulation, which results in considerable CO₂ emissions. In addition, peat oxidation causes smaller amounts of nitrous oxide (N₂O) and CH₄ emissions (Hiraishi *et al* 2014, Wilson

1
2
3 *et al* 2016). On top of that, ditches in drained peatlands carry larger amounts of dissolved
4 organic carbon (DOC) out of the peatland than in their natural state, and most of the DOC is
5 oxidized and released as CO₂ (Hiraishi *et al* 2014). In terms of global warming potential
6 (GWP) CO₂ is by far the most important GHG from degraded peatlands (Barthelmes *et al*
7 2015, Hiraishi *et al* 2014). The process of complete peat oxidation may take centuries for
8 deep peat deposits (Leifeld and Menichetti 2018). Therefore, each unit of drained peatland
9 is a contributor to global warming until all the peat is eventually decomposed. Besides the
10 protection of intact peatlands, an option to reverse this trend is the restoration of drained
11 peatlands through rewetting (Barthelmes *et al* 2015, Wilson *et al* 2013, Jaenicke *et al* 2010).
12 Similar to intact peatlands, rewetted peatlands absorb CO₂ from the atmosphere but emit
13 on average 46% more methane (Abdalla *et al* 2016). Net GHG emissions from rewetted
14 peatlands are in many cases much lower compared to drained peatlands, but the emission
15 reductions are variable over time and in different climatic regions and therefore not easily
16 generalizable (Barthelmes *et al* 2015, Wilson *et al* 2016). According to the most up-to-date
17 emission factors, rewetted peatlands over mid-term have higher GHG emissions than intact
18 peatlands (Barthelmes *et al* 2015, Wilson *et al* 2016, Hiraishi *et al* 2014). But despite
19 rewetted peatlands may remain 'negative' for the climate in an absolute sense, the GHG
20 mitigation compared to the previous drained status remains considerable.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

For climate policy-making in line with the Paris Agreement it is therefore crucial that emissions from degraded peatlands, which can be estimated with the help of IPCC Guidelines (Hiraishi *et al* 2014), are accounted for in mitigation pathways towards 1.5°C or 2°C in 2100. To our knowledge, which is based on the IPCC special report on 1.5° (Rogelj *et al* 2018), the IPCC special report on climate change and land (Smith *et al* 2019, Table 6.76), and the IAMC 1.5°C scenario explorer (Huppmann *et al* 2018), there is currently no single

1
2
3 integrated assessment modelling (IAM) study, which specifically accounts for the challenges
4 of peatland protection and restoration related mitigation and food security. Considering
5 emissions from degraded peatlands in mitigation pathways is first of all crucial due to the
6 legacy of historic peatland conversion. On top of that, most pathways consistent with 1.5°C
7 or 2°C include land-demanding mitigation options, such as bioenergy and afforestation, with
8 substantial consequences for the land system (Clarke *et al* 2014, Rogelj *et al* 2018).
9
10 However, it remains unclear so far if and to what extent land-demanding mitigation options
11 could negatively affect intact peatlands by clearance and drainage for energy crops in the
12 future. In this study, we estimate for the first time the potential GHG consequences of
13 peatland degradation in a likely 2°C mitigation pathway, along with increasing demand for
14 food and feed for a growing population. Moreover, we explore, also for the first time, the
15 potential of peatland protection and restoration policies for lowering GHG emissions from
16 degraded peatlands, including effects on food security, in a likely 2°C mitigation pathway.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 36 Methods

37
38 For this study, we developed a peatland module for the *MAGPIE 4 open-source*
39 *framework for modeling global land systems* (Dietrich *et al* 2019). The model version
40 including the peatland module we use here is MAGPIE 4.2.1 (Dietrich *et al* 2020) (see data
41 availability statement at the end of the article for details). MAGPIE combines economic and
42 biophysical approaches to simulate spatially explicit global scenarios of land use within the
43 21st century and the respective interactions with the environment (Figure S2-S3). The
44 MAGPIE framework has been used to simulate mitigation pathways for different Shared
45 Socioeconomic Pathways (SSPs) (Popp *et al* 2017) and contributed to several IPCC reports
46 (Rogelj *et al* 2018, Smith *et al* 2019). In particular, the MAGPIE framework has been used for
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 several studies on the environmental effects of terrestrial climate change mitigation
4
5 measures, including large-scale bioenergy production and afforestation (Humpenöder *et al*
6
7 2018, 2014, Popp *et al* 2014, Bonsch *et al* 2016, Popp *et al* 2011, Kreidenweis *et al* 2016). In
8
9 line with these earlier studies, we do not account for impacts of climate change on crop
10
11 yields and vegetation carbon stocks in this study. In the following we describe how we
12
13 initialize the peatland module, how we model future peatland dynamics and associated
14
15 GHG emissions, and how we implemented the peatland protection and restoration
16
17 scenarios. The SI provides a schematic overview of the study setup (Figure S1) and a
18
19 description of the main characteristic of the MAgPIE 4 modelling framework.
20
21
22
23
24
25
26
27

28 Initial peatland map

29 The peatland module is initialized with a map of present-day degraded and intact
30
31 peatlands (Figure S4), which we generated by combining a map of potential peatland area,
32
33 provided by the authors of Leifeld and Menichetti (2018) and Yu *et al* (2010), with country
34
35 level data on the status and extent of peatlands (intact and degraded) for the year 2015
36
37 from the Global Peatland Database (GPD) (Greifswald Mire Centre 2015). The map of
38
39 potential peatland distribution is mainly based on geomorphology and hydrology. On the
40
41 global level the area of the potential peatland map is about twice as large (~997 Mha)
42
43 compared to the country level peatland area data (~497 Mha). We downscaled the country
44
45 level peatland area (intact and degraded) to 0.5-degree resolution grid cell level by using the
46
47 potential peatland map as weight and upper limit. In a first step, the degraded and intact
48
49 peatland area of each country from GPD is distributed to the corresponding grid cells
50
51 weighted by the potential peatland map. In a second step, we limit the peatland area in
52
53 each grid cell to the area of the potential peatland map (for more details see source code in
54
55
56
57
58
59
60

1
2
3 supplementary data). Since there is some mismatch between GPD and the potential
4
5 peatland map, the area of the resulting map is slightly smaller than the area of GPD (~421
6
7 Mha vs. ~497 Mha globally). On the global level, degraded peatland in our map is about 5
8
9 Mha smaller compared to GPD, while intact peatland is about 71 Mha smaller. The source
10
11 code for compiling the peatland map is available as supplementary data.
12
13
14

15
16 Our spatially explicit map for the year 2015 provides for each grid cell the extent of
17
18 intact and degraded peatlands, but does not provide information on how degraded
19
20 peatland is used (for instance as cropland). The IPCC wetland GHG emission factors,
21
22 however, are differentiated into the land-use categories cropland, pasture and forestry (see
23
24 Table S1). Since MAgPIE has detailed land use information for each cell, we allocate
25
26 degraded peatland area in consecutive order to the land-use categories cropland, pasture
27
28 and forestry, which is comparable to the allocation rules used in Leifeld and Menichetti
29
30 (2018). To make sure that the initial distribution of degraded peatland to cropland, pasture
31
32 and forestry is consistent with our approach for modelling future peatland dynamics (see
33
34 next subsection), we account for the ratio of total peatland area to total land area in each
35
36 cell (hereafter referred to as scaling factor) in the initial distribution. In each of the three
37
38 steps (cropland, pasture, forestry), the allocation of degraded peatland to managed land is
39
40 constrained by the respective level of land use in MAgPIE in 2015 multiplied with the scaling
41
42 factor (see model code for details). Any unallocated degraded peatland area after these
43
44 three steps is assigned to an *unused* peatland category with area-weighted distribution
45
46 among cropland, pasture and forestry.
47
48
49
50
51
52
53

54 [Modelling future peatland dynamics](#)

55
56 Starting from this initial peatland map for the year 2015, we estimate future
57
58 peatland dynamics based on projections of changes in cropland, pasture and forestry from
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MAGPIE. We assume that changes of cropland, pasture and forestry translate into changes of degraded peatland proportional to the ratio of total peatland area to total land area in each cell (scaling factor between 0 and 1). Furthermore, we assume that the sum of the peatland categories intact, degraded, unused and rewetted remains constant over time in each cell, which also implies that the scaling factor is constant over time. The following example illustrates the mechanism used for projecting peatland dynamics (see equations in model code for details): In a given cell, the total land area is 50 Mha and the total peatland area is 10 Mha. Therefore, the scaling factor is 0.2 (10 Mha divided by 50 Mha). If cropland expands by 5 Mha, 1 Mha of intact peatland is converted to degraded peatland ($5 \text{ Mha} \times 0.2$). By multiplying changes in managed land with the scaling factor we implicitly assume that intact peatlands are distributed equally within a cell. If the totality of the cell would become cropland, degraded peatland would equal to the total peatland area ($50 \text{ Mha} \times 0.2 = 10 \text{ Mha}$). Expansion of cropland, pasture or forestry in cells with peatland does not necessarily result in degradation of intact peatlands. If degraded peatland in the *unused* category exists, the model will use this land first before converting intact peatland. To avoid double accounting of costs, we don't assume specific costs for the drainage of intact peatlands because MAGPIE already accounts for one-time costs of 8000 USD ha⁻¹ (2005 market exchange rate; MER) for the conversion of unmanaged to managed land.

If cropland, pasture or forestry decline, the process just works the other way around, i.e. degraded peatland is reduced proportional based on the scaling factor, with the difference that the reduction of degraded peatland results in an increase of the peatland categories *unused* or *rewetted*, depending on the pricing of GHG emissions from peatlands (see next sub-sections). The initial distribution of degraded peatland to cropland, pasture

1
2
3 and forestry makes sure that in case all managed land in a cell is abandoned for peatland
4 restoration, all degraded peatland will be rewetted.
5
6

7 8 GHG emission factors for degraded and rewetted peatlands 9

10 We use of the most up-to-date wetland GHG emission factors for drained and
11 rewetted peatlands (Table S1). The emission factors differ by emission type (CO₂, CH₄, N₂O,
12 CO₂-DOC), land type (cropland, pasture, forestry) and climate zone (tropical, temperate,
13 boreal). The GHG emission factors in Table S1 are taken from table 5 in Wilson *et al* (2016),
14 which is based on the 2013 Wetland Supplement to the 2006 IPCC Guidelines for National
15 Greenhouse Gas Inventories (Hiraishi *et al* 2014). In line with the IPCC Tier 1 approach we
16 use nutrient poor emission factors for boreal and nutrient rich emission factors for
17 temperate regions. Emission factors for tropical grassland have been added based on IPCC
18 (Hiraishi *et al* 2014). For the mapping of climate regions and cells we use the Koeppen-
19 Geiger climate classification (Table S2). GHG emissions from degraded peatlands are
20 obtained by multiplication of GHG emission factors for drained peatlands (t CO_{2eq} ha⁻¹ yr⁻¹)
21 with the respective area of degraded peatland (categories *degraded* and *unused*) in MAgPIE.
22 In line with IPCC guidelines, we assume the same GHG emission factors for all degraded
23 peatlands, irrespective of the point in time when the peat oxidation started. GHG emission
24 savings due to peatland restoration are obtained by multiplication of GHG emission
25 *reduction* factors (ER) for rewetted peatlands (difference between drained and rewetted
26 emission factors) with the respective area of rewetted peatland in MAgPIE. Impacts of
27 future climate change on emissions from peatlands are not accounted for in this study.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 54 Peatland protection and restoration 55

56 Peatland protection and restoration policies are implemented as economic process
57 in the model by pricing GHG emissions from degraded and rewetted peatlands (see section
58
59
60

1
2
3 below on scenario setup for details on the used CO₂ price path). Since the goal function of
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
restoration.

Scenario setup

We explore three main scenarios: *RCP2.6* (no peatland policy), *RCP2.6+PeatProt* (peatland protection policy) and *RCP2.6+PeatRestor* (peatland protection and restoration policy). These main scenarios are based on the Shared Socio-economic Pathway 2 (SSP2) (Popp *et al* 2017), which is a middle of the road scenario for future socio-economic developments, i.e. the main scenarios have the same assumptions for key parameters like future population, income, food demand, environmental protection and trade. However, how such key parameters develop in the future is uncertain - with potential impacts on peatland degradation or restoration. To assess this uncertainty, we run our main scenarios with alternative socio-economic assumptions (SSP1-5) (Figures S5-S6). Also, all scenarios share the same assumptions for land-based mitigation including large-scale production of 2nd generation bioenergy and CO₂ price path (Figures S7-S8), which are both taken from an existing RCP2.6 (likely 2°C in 2100) mitigation pathway available in the SSP database (IIASA 2018). We here use data from the "REMIND-MAgPIE - SSP2-26" pathway, which is *not* the so-called marker representation of SSP2-26. However, alternative representations of SSP2-26 are equally valuable and using the REMIND-MAgPIE (MAgPIE version 3) representation of SSP2-26 is the most consistent setup for this study since we here use a follow-up version of MAgPIE (version 4.2.1). In this study, bioenergy demand is an exogenous input to the model. Hence, the model accounts for emissions from land-use change (e.g. from deforestation) but does not account for CO₂ benefits resulting from the use of bioenergy. In all scenarios, the GHG price is applied on CO₂ emissions from the conversion of unmanaged land (forest and non-forest natural vegetation), as well as on CH₄ and N₂O emission from agricultural production (converted using GWP factors of 28 for CH₄ and 265 for N₂O). The pricing of land-related GHG emissions creates an economic incentive in the model to reduce these

emissions, e.g. by less conversion of unmanaged to managed land or by investments in emission-friendly technologies (implemented via marginal abatement cost curves) (Popp *et al* 2010). In the *RCP2.6* scenario, GHG emissions from the conversion of intact peatlands and historically degraded peatlands are not priced, i.e. these emissions are not considered in the decision-making process. However, GHG emissions from degraded peatlands are tracked by the model. In the *RCP2.6+PeatProt* scenario, the GHG price is applied on GHG emissions from degraded peatlands, which provides an incentive to stop further degradation of intact peatlands. In the *RCP2.6+PeatRestor* scenario, GHG emission from degraded peatlands are priced as in *RCP2.6+PeatProt*, and in addition the rewetting of degraded peatlands, which reduces GHG emissions compared to the drained state (Table S1), is rewarded based on the GHG price.

Results

General land dynamics

The results for our three main scenarios are similar in terms of global dynamics for cropland, pasture and forest throughout the 21st century (Figure 1). All scenarios show a strong increase of cropland for bioenergy production (412-447 Mha globally until 2100), and a smaller increase of forest area due to afforestation (134-149 Mha globally until 2100). These land expansions mainly come at the cost of other natural land and cropland for food production. Since demand for food, due to growing population and income, as well as demand for bioenergy, due to climate policy, are increasing throughout the 21st century (Figures S5-S7), reduction of cropland for food production is only possible under continued yield increases (Figure S12). Also, the results in terms of CO₂ emissions from changes in vegetation (e.g. from loss of natural land and afforestation) are similar for all scenarios

(Figure 3). CO₂ emissions from changes in vegetation are positive until around 2030 due to land conversion. After 2030, the CO₂ price applied on these emissions starts to increase rapidly (Figure S8), which alters land conversion patterns such that associated CO₂ emissions are strongly reduced. Moreover, global CO₂ emissions from changes in vegetation turn net-negative after 2030 due to afforestation based on existing national policies. In the *RCP2.6+PeatRestor* scenario, reduced conversion of other natural land results in more negative CO₂ emissions due to ongoing regrowth of vegetation (Figures 1 and 3).

No peatland policy

With respect to peatland area, all scenarios start in 2015 with degraded peatland area of 45.8 Mha and intact peatland area of 375.2 Mha globally (Table 1a, Figure 2). 36% of all degraded peatlands are located in the tropics (mainly Southeast Asia) and 47% are located in the boreal climate zone (mainly Europe and Russia) (Figures S4 and S14). In 2015, annual GHG emissions from degraded peatlands (CO₂, CH₄ and N₂O) including off-site CO₂ emissions from DOC amount to 1.50 Gt CO_{2eq} yr⁻¹ of which 82% (1.24 Gt CO₂ yr⁻¹) come from peat oxidation (Table 1b, Figure 4).

In the scenario without peatland policies (*RCP2.6*), global degraded peatland area increases between 2015 and 2100 by 10.3 Mha at the cost of intact peatlands, which corresponds to a relative increase of 22% compared to 2015 (Figure 2b). The strongest increase takes place in the tropics (5.7 Mha by 2100). The pace of peatland degradation throughout the 21st century partly coincides with the strong expansion of cropland for bioenergy production from about 2040 onwards (Figure 1). In consequence, annual global CO₂ emissions from degraded peatlands increase to 1.58 Gt CO₂ yr⁻¹ in 2100, which is about 25% higher compared to 1.24 Gt CO₂ yr⁻¹ in 2015 (Figure 3). In addition, also other emissions from degraded peatlands (DOC, CH₄ and N₂O) increase by about 25% globally between 2015

1
2
3 and 2100, resulting in total emission from degraded peatlands of 1.92 Gt CO_{2eq} yr⁻¹ in 2100
4
5 (Figure 4). Due to the CO₂ emissions from degraded peatlands, global net CO₂ emissions
6
7 from the land system including a small sink due to afforestation remain net-positive
8
9 throughout the 21st century (1.23 Gt CO₂ yr⁻¹ in 2100) (Figure 3).
10
11

12 13 Peatland protection policy

14
15 The scenario with peatland protection through pricing of GHG emissions from
16
17 degraded peatlands (*RCP2.6+PeatProt*) shows substantially less degradation of intact
18
19 peatlands throughout the 21st century (Figure 2b), resulting in lower GHG emission from
20
21 degraded peatlands (Figure 4), despite the same considerable expansion of cropland for
22
23 bioenergy production (Figure 1). By 2100, degraded peatlands increase only by 2.3 Mha
24
25 globally, compared to 10.3 Mha in the scenario without peatland policy (*RCP2.6*). About 50%
26
27 of the degradation happens until 2030, which coincides with the strong rise of the CO₂ price
28
29 from 2030 onwards (Figure S8). In our price path, the global CO₂ price in 2030 is 39 USD
30
31 tCO₂⁻¹. In consequence, emissions from degraded peatlands increase only slightly until 2030,
32
33 and remain rather constant later on (1.59 Gt CO_{2eq} yr⁻¹ in 2100, Figure 4). Notably, most of
34
35 the peatland emissions in this scenario originate from historic peatland degradation. The
36
37 global net CO₂ emissions from the land system are 0.96 Gt CO₂ yr⁻¹ in 2100 (Figure 3), which
38
39 is lower compared to 1.23 Gt CO₂ yr⁻¹ in the scenario without peatland policy. However,
40
41 even in the case of peatland protection, the land system remains a net source of CO₂
42
43 throughout the 21st century, owing to the legacy of historic peatland conversion.
44
45
46
47
48
49
50

51 Peatland protection and restoration policy

52
53 Finally, the scenario with peatland protection *and* restoration policies
54
55 (*RCP2.6+PeatRestor*) shows substantial rewetting of historically degraded peatlands next to
56
57 the protection of intact peatlands in the future (Figure 2b) despite strong expansion of
58
59
60

1
2
3 cropland for bioenergy production (Figure 1). Between 2015 and 2100 global degraded
4 peatland area shrinks by 29.4 Mha due to rewetting, which corresponds to a relative
5 reduction of about 60% compared to 2015. Notably, about 50% of the global peatland
6 restoration (rewetting) occurs between 2025 and 2035 (Figure 2b), which indicates that
7 peatland restoration is a cost-efficient mitigation measure at relatively low CO₂ prices (22
8 USD tCO₂⁻¹ in 2025, Figure S8). About half of the peatland rewetting takes place in the
9 tropical (Southeast Asia) and the boreal climate zone (Europe and Russia), respectively
10 (Figure S14). In consequence, CO₂ emissions from degraded peatlands decrease
11 considerably from 2025 onwards to 0.36 Gt CO₂ yr⁻¹ in 2100 (Figure 4). At the same time,
12 CH₄ emissions more than double between 2015 and 2100 due to considerable rewetting of
13 degraded peatlands. However, the net effect is a reduction of emissions from peatlands to
14 0.61 Gt CO_{2eq} yr⁻¹ in 2100, which is qualitatively in line with Günther et al (2020). Because of
15 the much lower CO₂ emissions from peatlands, together with the slightly higher carbon sink
16 due to vegetation regrowth, the land system turns into a global net carbon sink by 2075 in
17 this scenario (Figure 3).

40 Socio-economic impacts and uncertainty

41
42 Future peatland degradation and restoration is small in terms of area compared to
43 overall land-use change, in particular compared land expansion for bioenergy production
44 (e.g. 32 Mha restored peatland vs. > 400 Mha bioenergy area globally in 2100 in
45 *RCP2.6+PeatRestor*, Figures 1 and 2). However, despite the relatively small area changes,
46 peatland protection and restoration measures could have economic impacts because they
47 imply that expansion of managed land is to some extent constrained, which could increase
48 the overall costs of agricultural production. In our peatland protection and restoration
49 scenario (*RCP2.6+PeatRestor*), about 50% of the global peatland restoration happens until
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2035 (Figure 2b). As proxy for the near-term costs of peatland protection and restoration
4
5 measures, we compare the change of total agricultural production costs by 2035 (index
6
7 relative to 2015) between our scenarios. Total agricultural production costs include the
8
9 direct costs for peatland restoration as well as costs for labour, capital, fertilizer,
10
11 technological change, intraregional transport, land conversion and GHG emissions. Our
12
13 results show that peatland protection and restoration measures (*RCP2.6+PeatProt*,
14
15 *RCP2.6+PeatRestor*) hardly increase the whole system costs on top of costs for agricultural
16
17 production including climate policy (*RCP2.6*) at the level of geopolitical world regions (Figure
18
19 5). As proxy for near-term food security we use a food price index (Paasche price index). The
20
21 food prices used for calculating the food price index are shadow prices, which reflect the
22
23 marginal costs of one additional unit of production (Humpenöder *et al* 2018). Similar to the
24
25 agricultural production costs, food prices in 2035 show almost no reaction to peatland
26
27 protection and restoration policies on top of climate policy for all world regions. This result
28
29 holds true also for the longer-term projection of global food prices until 2100 (Figure S10).
30
31 Moreover, food availability, which is the outcome of a price-elastic food demand response
32
33 function, is identical across all three scenarios at global level (Figure S11). In addition, trade-
34
35 patterns between the regions are not altered by peatland protection and restoration
36
37 policies (Figure S13).

38
39
40
41
42
43
44
45
46
47 In our SSP2-based main scenarios, peatland protection and restoration measures
48
49 substantially reduce degraded peatland without major increases in production costs and
50
51 food prices. Qualitatively, this finding is robust under alternative socio-economic
52
53 assumptions (SSP1-5). Under all SSPs, peatland protection and restoration measures reduce
54
55 degraded peatland, while agricultural production costs and food prices remain rather
56
57 unchanged (Figure 5). However, there is some uncertainty regarding the level of peatland
58
59

1
2
3 degradation and associated costs, depending on the underlying socio-economic
4 assumptions (e.g. in Sub-Saharan Africa). For peatland restoration, we additionally tested
5 the robustness of our results against lower and higher restoration costs compared to our
6 default assumptions (see methods). Overall, the sensitivity is small (Figure S9). A noticeable
7 difference in restoration area occurs in the boreal climate zone around 2060 (8.2, 10.1 and
8 12.0 Mha for high, medium and low restoration costs, respectively).
9
10
11
12
13
14
15
16
17
18

19 Discussion

20 Estimates of peatland area and GHG emissions

21
22 Our estimate for global present-day degraded peatland area and associated GHG
23 emissions is comparable to other studies (Table 1). However, there are differences at the
24 level of climate zones, which are only reported by two of these studies (Leifeld and
25 Menichetti 2018, Leifeld *et al* 2019). Our estimate for GHG emissions in the tropical climate
26 zone is lower, which is consistent with the smaller degraded peatland area in our data set.
27 Moreover, we disregard emissions from peat extraction, which further explains why our
28 estimate for GHG emissions in the tropical zone is lower. On the other hand, our estimate
29 for GHG emissions in the temperate and boreal climate zone is higher, while the
30 corresponding degraded area is similar. This indicates that distribution of degraded peatland
31 to cropland, pasture and forestry, which partly have diverging emission factors (Table S1), is
32 different. Although we use similar rules for the distribution of degraded peatland to
33 managed land as Leifeld and Menichetti (2018), different land-use data sets have been used
34 for this allocation.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54
55 According to our scenario results, protection of intact peatlands and restoration of
56 degraded peatlands considerably reduce CO₂ emissions from peat degradation at low costs.
57
58
59
60

1
2
3 This finding is in line with case studies from Indonesia (Jaenicke *et al* 2010) and Ireland
4
5 (Wilson *et al* 2013). Moreover, peatland restoration has developed over the last 25 years
6
7 from small-scale to landscape level restoration (Chimner *et al* 2017), which underpins the
8
9 plausibility of our scenario results. However, the success of peatland restoration depends on
10
11 several factors including the time elapsed since the end of peatland use, the starting
12
13 conditions and the restoration techniques (Höper *et al* 2008). In line with previous studies,
14
15 our restoration scenario shows higher CH₄ emissions due to anaerobic conditions caused by
16
17 rewetting (Abdalla *et al* 2016). Finally, we find that the net effect of substantially lower CO₂
18
19 emissions and slightly higher CH₄ emissions, in terms of GWP, is a net reduction of GHG
20
21 emissions from peatlands following restoration, which is supported by a recently published
22
23 study (Günther *et al* 2020).
24
25
26
27
28

29 30 Modelling peatland dynamics

31
32 For estimating future peatland degradation and rewetting, we multiply changes in
33
34 managed land with a peatland scaling factor (proportion of total peatland area over total
35
36 land area). With this approach we implicitly assume that intact peatlands are distributed
37
38 equally within a cell. We acknowledge that this is a simplification, which might result in
39
40 under- or overestimation of future peatland degradation. However, in the absence of better
41
42 data on the spatial distribution of intact peatlands, we consider our approach as appropriate
43
44 since it is plausible (equal distribution) and allows to compare quantitatively various
45
46 scenarios of future peatland dynamics in a meaningful way. Given our assumptions, all
47
48 intact peatland in a cell will be degraded if all available land in this cell is used for
49
50 production. In the contrary case, when all managed land in a cell is abandoned for peatland
51
52 restoration, all degraded peatland will be rewetted. These two boundary cases define the
53
54 range of possible future peatland dynamics within our approach.
55
56
57
58
59
60

1
2
3 In line with IPCC guidelines, we assume the same GHG emission factors for all
4 degraded peatlands, irrespective of the point in time when the peat oxidation started. For
5 deep peat deposits, the process of complete peat oxidation may indeed take centuries
6 (Leifeld and Menichetti 2018), but shallow peat layers (and more so in case of fires) may be
7 close to depletion already after 5 years (Hooijer *et al* 2012). In any case where the peat
8 deposit is close to depletion emission rates would gradually slow down (following kinetics
9 more similar to mineral soils) rapidly approaching zero net emissions. Therefore, our results
10 might overestimate GHG emissions from degraded peatlands to some extent. On the other
11 hand, we here a) do not account for impacts of climate change (e.g. changes in temperature
12 and precipitation), which might reduce the carbon storage potential of peatlands in the
13 future (Leng *et al* 2019), and b) disregard GHG emissions related to peat extraction.

30 Limitations of our global model-based scenario projections

32 Our scenario projections until 2100 must not be interpreted as exact predictions.
33 Instead, our land-use projections including peatland dynamics reflect plausible future
34 trajectories given the underlying socio-economic and climate policy assumptions. The
35 numerical modelling makes these trajectories comparable with each other. The scope of our
36 spatially-explicit approach is modelling future land-use change including peatland dynamics
37 driven by demand for food and land-based mitigation with global coverage. Since peatlands
38 cover only about 3% of the total land area, also projected changes of peatland area are
39 small compared to changes in major land-use types such cropland or pasture (see Figures 1
40 and 2). Due to the small-scale nature of peatland dynamics and because demand for food
41 and bioenergy is defined at the level of 12 geopolitical regions there is considerable
42 flexibility within each region for adjustment of production patterns in the case of peatland
43 protection and/or restoration policies. In our results, peatland protection and restoration

1
2
3 policies have almost no economic impact in terms of implementation costs and food
4 security at regional level (Figure 5). In reality, however, agriculture on drained peatlands
5 may in some local contexts play an important role for food security due to prevailing
6 subsistence agriculture, e.g. in low income countries in Sub-Saharan Africa. Our approach is
7 not capable of capturing such local effects due to its global scale scope.
8
9
10
11
12
13
14
15

16 Conclusion

17
18
19 Our results indicate that a climate policy with land-based mitigation but without
20 peatland protection and restoration measures could result in degradation of presently intact
21 peatlands, associated with an increase of GHG emissions from degraded peatlands in the
22 future. Moreover, even in the case of peatland protection, there are still considerable GHG
23 emissions from degraded peatlands, mostly owing to the historic conversion of intact
24 peatlands. Due to the emissions from degraded peatlands, the land system remains a
25 considerable net source of CO₂ throughout the 21st century globally in both cases (*RCP2.6*
26 and *RCP2.6+PeatProt*), despite negative emissions from afforestation. This result is in
27 contrast to the outcome of current mitigation pathways with land-based mitigation
28 measures, of which most do not include emissions from degraded peatlands, and therefore
29 report declining global CO₂ emissions from land system throughout the 21st century,
30 resulting in net-negative CO₂ emissions by 2100 in most scenarios (Huppmann *et al* 2018)
31 (Figure S15). Therefore, by omitting peatland emissions current mitigation pathways likely
32 overestimate the mitigation potential of the land system. However, in line with other
33 studies (Leifeld *et al* 2019), our results indicate that it is possible to reconcile land use and
34 GHG emissions in mitigation pathways through a peatland protection *and* restoration policy
35 (*RCP2.6+PeatRestor*). Our results suggest that the land system would turn into a global net
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 carbon sink by 2100, as projected by current mitigation pathways, if about 60% of present-
4
5 day degraded peatlands, mainly in the tropical and boreal climate zone, would be rewetted
6
7 in the coming decades, next to the protection of intact peatlands. Therefore, peatland
8
9 protection *and* restoration are key for climate change mitigation. At the same time, our
10
11 results indicate that the implementation costs of peatland protection and restoration
12
13 measures are low, and that there are almost no impacts on regional food security.
14
15
16
17
18

19 Acknowledgments

20
21 This work received funding from the NAVIGATE and the ENGAGE project of the European
22
23 Union's Horizon 2020 research and innovation program under grant agreements 821124
24
25 and 821471, respectively. The work of Kristine Karstens was funded by the DFG Priority
26
27 Program "Climate Engineering: Risks, Challenges, Opportunities?" (SPP 1689) and
28
29 specifically the CEMICS2 project (grant no. ED78/3-2).

30 Data availability statement

31
32 The data that support the findings of this study are openly available at
33
34 <http://doi.org/10.5281/zenodo.1418752>, which includes the source code (as developed on
35
36 <https://github.com/magpiemodel>) and input data for MAGPIE 4.2.1. The model
37
38 documentation can be found at <https://rse.pik-potsdam.de/doc/magpie/4.2.1/>.
39
40 The source code for compiling a spatially explicit map of intact and degraded peatlands, and
41
42 the source code for the figure plotting are available as supplementary data. The figures
43
44 were created with ggplot2 (Wickham 2016).

45 References

- 46 Abdalla M, Hastings A, Truu J, Espenberg M, Mander Ü and Smith P 2016 Emissions of
47 methane from northern peatlands: a review of management impacts and implications
48 for future management options *Ecol. Evol.* **6** 7080–102
- 49 Barthelmes A, Couwenberg J, Risager M, Tegetmeyer C and Joosten H 2015 *Peatlands and*
50 *Climate in a Ramsar context: a Nordic-Baltic Perspective. TemaNord 2015:544*
51 (Copenhagen K: Nordic Council of Ministers)
- 52 Bonsch M, Humpenöder F, Popp A, Bodirsky B, Dietrich J P, Rolinski S, Biewald A, Lotze-
53 Campen H, Weindl I, Gerten D and Stevanovic M 2016 Trade-offs between land and
54 water requirements for large-scale bioenergy production *GCB Bioenergy* **8** 11–24
- 55 Chimner R A, Cooper D J, Wurster F C and Rochefort L 2017 An overview of peatland
56 restoration in North America: where are we after 25 years?: Peatland restoration in
57 North America *Restor. Ecol.* **25** 283–92

- Clarke L, Jiang K, Akimoto K, Babiker M, Blanford G, Fisher-Vanden K, Hourcade J-C, Krey V, Kriegler E, Löschel A, McCollum D, Paltsev S, Rose S, Shukla P R, Tavoni M, van der Zwaan B and van Vuuren D P 2014 Assessing Transformation Pathways *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and J.C. Minx (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Dietrich J P, Bodirsky B L, Humpenöder F, Weindl I, Stevanović M, Karstens K, Kreidenweis U, Wang X, Mishra A, Klein D, Ambrósio G, Araujo E, Yalew A W, Baumstark L, Wirth S, Giannousakis A, Beier F, Chen D M-C, Lotze-Campen H and Popp A 2019 MAGPIE 4 – a modular open-source framework for modeling global land systems *Geosci. Model Dev.* **12** 1299–317
- Dietrich J P, Bodirsky B L, Weindl I, Humpenöder F, Stevanovic M, Kreidenweis U, Wang X, Karstens K, Mishra A, Beier F D, Bacca E J M, Klein D, Ambrósio G, Araujo E, Biewald A, Lotze-Campen H and Popp A 2020 *MagPIE - An Open Source land-use modeling framework - Version 4.2.1* (doi: 10.5281/zenodo.1418752) Online: <https://github.com/magpiemodel/magpie>
- Edenhofer O, Pichs-Madruga R, Sokona Y, Kadner S, Minx J C, Brunner S, Agrawala S, Baiocchi G, Bashmakov I A, Blanco G, Broome J, Bruckner T, Bustamante M, Clarke L, Grand M C, Creutzig F, Cruz-Núñez X, Dhakal S, Dubash N K, Eickemeier P, Farahani E, Fishedick M, Fleurbaey M, Gerlagh R, Gómez-Echeverri L, Gupta S, Harnisch J, Jiang K, Jotzo F, Kartha S, Klasen S, Kolstad C, Krey V, Kunreuther H, Lucon O, Masera O, Mulugetta Y, Norgaard R B, Patt A, Ravindranath N H, Riahi K, Roy J, Sagar A, Schaeffer R, Schlömer S, Seto K C, Seyboth K, Sims R, Smith P, Somanathan E, Stavins R, Stechow C von, Sterner T, Sugiyama T, Suh S, Ürges-Vorsatz D, Urama K, Venables A, Victor D G, Weber E, Zhou D, Zou J and Zwickel T 2014 Technical Summary *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed O Edenhofer, R Pichs-Madruga, Y Sokona, E Farahani, S Kadner, K Seyboth, A Adler, I Baum, S Brunner, P Eickemeier, B Kriemann, J Savolainen, S Schlömer, C von Stechow, T Zwickel and J C Minx (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press)
- Glenk K and Martin-Ortega J 2018 The economics of peatland restoration *J. Environ. Econ. Policy* **7** 345–62
- Greifswald Mire Centre 2015 *Global Peatland Database* (Greifswald, Germany: Greifswald Mire Centre) Online: <https://greifswaldmoor.de/global-peatland-database-en.html>
- Günther A, Barthelmes A, Huth V, Joosten H, Jurasinski G, Koebisch F and Couwenberg J 2020 Prompt rewetting of drained peatlands reduces climate warming despite methane emissions *Nat. Commun.* **11** 1–5

- 1
2
3 Hiraishi T, Krug T, Tanabe K, Srivastava N, Baasansuren J, Fukuda M and Troxler T G 2014
4 *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas*
5 *Inventories: Wetlands* (Switzerland: IPCC) Online: [http://www.ipcc-](http://www.ipcc-nggip.iges.or.jp/public/wetlands/)
6 [nggip.iges.or.jp/public/wetlands/](http://www.ipcc-nggip.iges.or.jp/public/wetlands/)
7
8
9 Hooijer A, Page S, Canadell J G, Silvius M, Kwadijk J, Wösten H and Jauhiainen J 2010
10 Current and future CO₂ emissions from drained peatlands in Southeast Asia
11 *Biogeosciences* **7** 1505–14
12
13 Hooijer A, Page S, Jauhiainen J, Lee W A, Lu X X, Idris A and Anshari G 2012 Subsidence
14 and carbon loss in drained tropical peatlands *Biogeosciences* **9** 1053–71
15
16 Höper H, Augustin J, Cagampan J P, Drösler M, Lundin L, Moors E, Vasander H,
17 Waddington J M and Wilson D 2008 Restoration of peatlands and greenhouse gas
18 balances *Peatlands and climate change* ed edited by Maria Strack (International:
19 International Peat Society) pp 182–210
20
21
22 Humpenöder F, Popp A, Bodirsky B L, Weindl I, Biewald A, Hermann Lotze-Campen,
23 Dietrich J P, Klein D, Kreidenweis U, Müller C, Susanne Rolinski and Stevanovic M
24 2018 Large-scale bioenergy production: how to resolve sustainability trade-offs?
25 *Environ. Res. Lett.* **13** 024011
26
27
28 Humpenöder F, Popp A, Dietrich J P, Klein D, Lotze-Campen H, Bonsch M, Bodirsky B L,
29 Weindl I, Stevanovic M and Müller C 2014 Investigating afforestation and bioenergy
30 CCS as climate change mitigation strategies *Environ. Res. Lett.* **9** 064029
31
32
33 Huppmann D, Kriegler E, Krey V, Riahi K, Rogelj J, Rose S K, Weyant J, Bauer N, Bertram
34 C, Bosetti V, Calvin K, Doelman J, Drouet L, Emmerling J, Frank S, Fujimori S,
35 Gernaat D, Grubler A, Guivarch C, Haigh M, Holz C, Iyer G, Kato E, Keramidas K,
36 Kitous A, Leblanc F, Liu J-Y, Löffler K, Luderer G, Marcucci A, McCollum D,
37 Mima S, Popp A, Sands R D, Sano F, Strefler J, Tsutsui J, Van Vuuren D, Vrontisi Z,
38 Wise M and Zhang R 2018 *IAMC 1.5°C Scenario Explorer and Data hosted by IIASA*
39 *(doi: 10.22022/SR15/08-2018.15429)* (Integrated Assessment Modeling Consortium
40 & International Institute for Applied Systems Analysis) Online:
41 <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer>
42
43
44 IIASA 2018 *SSP Database (version 2.0)* (Laxenburg: International Institute for Applied
45 Systems Analysis) Online: <https://secure.iiasa.ac.at/web-apps/ene/SspDb>
46
47
48 Jaenicke J, Wösten H, Budiman A and Siebert F 2010 Planning hydrological restoration of
49 peatlands in Indonesia to mitigate carbon dioxide emissions *Mitig. Adapt. Strateg.*
50 *Glob. Change* **15** 223–39
51
52
53 Jauhiainen J, Page S E and Vasander H 2016 Greenhouse gas dynamics in degraded and
54 restored tropical peatlands *Mires Peat* 1–12
55
56
57 Joosten H 2010 *The Global Peatland CO₂ Picture: Peatland Status and Drainage Related*
58 *Emissions in All Countries of the World* (Ede, The Netherlands: Wetland
59 International)
60

- 1
2
3 Joosten H, Sirin A, Couwenberg J, Laine J and Smith P 2016 The role of peatlands in climate
4 regulation *Peatland Restoration and Ecosystem Services: Science, Policy and*
5 *Practice* Ecological Reviews ed A Bonn, H Joosten, M Evans, R Stoneman and T
6 Allott (Cambridge: Cambridge University Press) pp 63–76
7
8
9 Kreidenweis U, Humpenöder F, Stevanović M, Bodirsky B L, Elmar Kriegler, Lotze-Campen
10 H and Popp A 2016 Afforestation to mitigate climate change: impacts on food prices
11 under consideration of albedo effects *Environ. Res. Lett.* **11** 085001
12
13 Leifeld J and Menichetti L 2018 The underappreciated potential of peatlands in global
14 climate change mitigation strategies *Nat. Commun.* **9** 1071
15
16 Leifeld J, Wüst-Galley C and Page S 2019 Intact and managed peatland soils as a source and
17 sink of GHGs from 1850 to 2100 *Nat. Clim. Change* 1–3
18
19 Leng L Y, Ahmed O H and Jalloh M B 2019 Brief review on climate change and tropical
20 peatlands *Geosci. Front.* **10** 373–80
21
22
23 Levy P E and Gray A 2015 Greenhouse gas balance of a semi-natural peatbog in northern
24 Scotland *Environ. Res. Lett.* **10** 094019
25
26 Moxey A and Moran D 2014 UK peatland restoration: Some economic arithmetic *Sci. Total*
27 *Environ.* **484** 114–20
28
29
30 Popp A, Calvin K, Fujimori S, Havlik P, Humpenöder F, Stehfest E, Bodirsky B L, Dietrich J
31 P, Doelmann J C, Gusti M, Hasegawa T, Kyle P, Obersteiner M, Tabeau A,
32 Takahashi K, Valin H, Waldhoff S, Weindl I, Wise M, Kriegler E, Lotze-Campen H,
33 Fricko O, Riahi K and Vuuren D P van 2017 Land-use futures in the shared socio-
34 economic pathways *Glob. Environ. Change* **42** 331–45
35
36
37 Popp A, Dietrich J P, Lotze-Campen H, Klein D, Bauer N, Krause M, Beringer T, Gerten D
38 and Edenhofer O 2011 The economic potential of bioenergy for climate change
39 mitigation with special attention given to implications for the land system *Environ.*
40 *Res. Lett.* **6** 034017
41
42
43 Popp A, Lotze-Campen H and Bodirsky B 2010 Food consumption, diet shifts and associated
44 non-CO₂ greenhouse gases from agricultural production *Glob. Environ. Change* **20**
45 451–62
46
47 Popp A, Rose S K, Calvin K, van Vuuren D P, Dietrich J P, Wise M, Stehfest E,
48 Humpenöder F, Kyle P, Vliet J V, Bauer N, Lotze-Campen H, Klein D and Kriegler E
49 2014 Land-use transition for bioenergy and climate stabilization: model comparison
50 of drivers, impacts and interactions with other land use based mitigation options *Clim.*
51 *Change* **123** 495–509
52
53
54 Rogelj J, Shindell D, Jiang K, Fifita S, Forster P, Ginzburg V, Handa C, Khesghi H,
55 Kobayashi S, Kriegler E, Mundaca L, Séférian R and Vilariño M V 2018 Mitigation
56 pathways compatible with 1.5°C in the context of sustainable development *Special*
57 *Report on the impacts of global warming of 1.5 °C* (Geneva: Intergovernmental Panel
58 on Climate Change) Online: <http://www.ipcc.ch/report/sr15/>
59
60

1
2
3 Smith P, Campbell D, Cherubini F, Grassi G, Nam V, Lwasa S, McElwee P, Saigusa N,
4 Soussana J-F, Angel M, Arias-Navarro C, Cavalett O, Cowie A, House J, Popp A,
5 Roe S, Slade R, Abdulla A, Noble I, Yamagata Y, Zatari T and Manning F 2019
6 Interlinkages between desertification, land degradation, food security and greenhouse
7 gas fluxes: Synergies, trade-offs and integrated response options *Climate Change and*
8 *Land: an IPCC special report on climate change, desertification, land degradation,*
9 *sustainable land management, food security, and greenhouse gas fluxes in terrestrial*
10 *ecosystems* (Geneva: Intergovernmental Panel on Climate Change) Online:
11 <https://www.ipcc.ch/srccl>

12
13
14 Wickham H 2016 *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New York)
15 Online: <https://ggplot2.tidyverse.org>

16
17
18 Wilson D, Blain D and Couwenberg J 2016 Greenhouse gas emission factors associated with
19 rewetting of organic soils *Mires Peat* **17** 4 28 pp

20
21 Wilson D, Müller C and Renou-Wilson F 2013 Carbon emissions and removals from Irish
22 peatlands: present trends and future mitigation measures *Ir. Geogr.* **46** 1–23

23
24 Yu Z, Loisel J, Brosseau D P, Beilman D W and Hunt S J 2010 Global peatland dynamics
25 since the Last Glacial Maximum *Geophys. Res. Lett.* **37**

26 27 28 29 30 Author Contributions Statement

31
32 F.H. expanded the MAgPIE modelling framework by a peatland module and applied it for
33 this study. F.H. designed the scenario setup, analysed the results, prepared all figures and
34 tables, and wrote the main manuscript text with important contributions from A.P and
35 H.L.C. K.K. contributed to the peatland module development. J.L, L.M. and A.B. provided
36 input data on extent and the state of peatlands. All authors reviewed the manuscript.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures and Tables

a) Degraded peatland area (Mha)					
	This study	(Leifeld and Menichetti 2018)	(Leifeld <i>et al</i> 2019)	(Joosten 2010)	
	<i>2015</i>	<i>2015</i>	<i>2015</i>	<i>2008</i>	
tropical	16.7	24.2	20.3		
temperate	7.6	10.6	26.7		
boreal	21.6	16.2	47.0		
global	45.8	50.9	47.0	42.6	

b) Annual GHG emissions from degraded peatlands (Gt CO_{2eq} yr⁻¹)					
	CO ₂	CO _{2eq}	CO _{2eq}	CO _{2eq}	CO ₂
tropical	0.76	0.88	1.48	1.26	
temperate	0.14	0.18	0.16	0.27	
boreal	0.33	0.44	0.26		
global	1.24	1.50	1.91	1.53	1.30

Table 1: Comparison of degraded peatland area and associated GHG emissions with other studies. For each study, the year for which the data is representative is provided in italic font. a) degraded peatland area in Mha. b) annual GHG emissions from degraded peatlands in Gt CO₂ yr⁻¹ or CO_{2eq} yr⁻¹ (depending on data availability).




Figure 1: Global change of major land use types (indicated by colour) over time until 2100 compared to 2015 for 3 scenarios (left, middle, right). Global land cover in 2015 amounts to 1511 Mha for cropland (food and bioenergy), 3260 Mha for pasture, 3989 Mha for forest and 4039 Mha for other land including urban areas (12805 Mha in total).




Figure 2: Peatland area in tropical, temperate and boreal climate zones. Panel a shows the status of degraded and intact peatlands in 2015. Panel b shows changes of peatland area over time until 2100 compared to 2015 for 3 scenarios (left, middle, right). The status of peatlands is indicated by colour, climate zones are indicated by corresponding colour gradients.




Figure 3: Land-related CO_2 emissions between 2015 and 2100 for 3 scenarios (left, middle, right) in tropical, temperate and boreal climate zones from two sources: change of vegetation carbon stocks (e.g. due to deforestation or afforestation) and peatland-related emissions (decomposition of peat in degraded/drained peatlands, accumulation of peat in rewetted peatlands). Please note that the CO_2 emissions shown here include only direct human effects. Indirect effects (e.g. due to climate change) are not accounted for. Emission sources are indicated by colour, climate zones are indicated by corresponding colour gradients.




Figure 4: Annual GHG emissions (CO_2 , CH_4 and N_2O) and off-site CO_2 emissions (DOC) from degraded and rewetted peatlands between 2015 and 2100 for 3 scenarios (left, middle, right) in tropical, temperate and boreal climate zones. Emission types are indicated by colour, climate zones are indicated by corresponding colour gradients.




Figure 5: Comparison of changes in degraded peatland area (left), agricultural production costs (middle) and food prices (right) for 8 geopolitical regions with at least 1 Mha of peatland (total peatland area in brackets; JPN, MEA and IND excluded) across 3 scenarios (indicated by color) for the year 2035. Changes are shown as index relative to 2015. Full color points indicate the results for the 3 SSP2-based main scenarios. Colored transparent bars show the corresponding robustness of these results under alternative socio-economic assumptions (SSP 1-5). For Sub-Saharan Africa (SSA) we illustratively label the minimum and maximum values of the uncertainty range. See Figure S3 for the definition of regions.

Supplementary Materials

Supplementary information is provided as additional PDF file. In addition, we provide the source code for the figure plotting and the source code for the compilation of the initial peatland map as supplementary data.